

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 June 2003 (05.06.2003)

PCT


(10) International Publication Number  
**WO 03/045937 A1**

- (51) International Patent Classification<sup>7</sup>: **C07D 401/12**, 211/18, 409/12, 401/04, 409/04, A61K 31/454, A61P 11/06
- (21) International Application Number: PCT/EP02/13218
- (22) International Filing Date:  
25 November 2002 (25.11.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/334,653 30 November 2001 (30.11.2001) US
- (71) Applicant: **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH).
- (72) Inventors: **DU BOIS, Daisy, Joe**; 1180 Welch Road, Palo Alto, CA 94304 (US). **KERTESZ, Denis, John**; 459 Victory Avenue, Mountain View, CA 94043 (US). **SJOGREN, Eric, Brian**; 442 Dell Avenue, Mountain View, CA 94043 (US). **SMITH, David, Bernard**; 218 West 40th Avenue, San Mateo, CA 94403 (US). **WANG, Beihan**; 3500 Granada Ave., Apt. 301, Santa Clara, CA 95051 (US).
- (74) Agent: **WAECHTER, Dieter**; Grenzacherstrasse 124, CH-4070 Basle (CH).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 03/045937 A1

(54) Title: N-UREIDO-PIPERIDINES AS ANTAGONISTS VIII FOR CCR-3 RECEPTOR



(57) Abstract: The invention relates to compounds of Formula (I), wherein R<sup>1</sup>, R<sup>5</sup>, A, L, and X are as defined in the specification. The compounds are useful as CCR-3 receptor antagonists, and therefore, may be used for treatment of CCR-3 mediated diseases.

## N-UREIDO-PIPERIDINES AS ANTAGONISTS VIII FOR CCR-3 RECEPTOR

This invention relates to piperidine derivatives that are CCR-3 receptor antagonists, pharmaceutical compositions containing them, their use, and methods for preparing them.

Tissue eosinophilia is a feature of a number of pathological conditions such as  
5 asthma, rhinitis, eczema and parasitic infections (see Bousquet, J. et al., N. Eng. J. Med.  
323: 1033-1039 (1990) and Kay, A. B. and Corrigan, C. J., Br. Med. Bull. 48:51-64 (1992)).  
In asthma, eosinophil accumulation and activation are associated with damage to  
bronchial epithelium and hyperresponsiveness to constrictor mediators. Chemokines such  
as RANTES, eotaxin and MCP-3 are known to activate eosinophils (see Baggiolini, M. and  
10 Dahinden, C. A., Immunol. Today. 15:127-133 (1994), Rot, A. M. et al., J. Exp. Med. 176,  
1489-1495 (1992) and Ponath, P. D. et al., J. Clin. Invest., Vol. 97, #3, 604-612 (1996)).  
However, unlike RANTES and MCP-3 which also induce the migration of other leukocyte  
cell types, eotaxin is selectively chemotactic for eosinophils (see Griffith-Johnson, D. A. et  
al., Biochem. Biophys. Res. Commun. 197:1167 (1993) and Jose, P. J. et al., Biochem.  
15 Biophys. Res. Commun. 207, 788 (1994)). Specific eosinophil accumulation was observed  
at the site of administration of eotaxin whether by intradermal or intraperitoneal injection  
or aerosol inhalation (see Griffith-Johnson, D. A. et al., Biochem. Biophys. Res. Commun.  
197:1167 (1993); Jose, P. J. et al., J. Exp. Med. 179, 881-887 (1994); Rothenberg, M. E. et  
al., J. Exp. Med. 181, 1211 (1995) and Ponath, P. D., J. Clin. Invest., Vol. 97, #3, 604-612  
20 (1996)).

Glucocorticoids such as dexamethasone, methprednisolone and hydrocortisone have  
been used for treating many eosinophil-related disorders, including bronchial asthma (R.  
P. Schleimer et al., Am. Rev. Respir. Dis., 141, 559 (1990)). The glucocorticoids are  
believed to inhibit IL-5 and IL-3 mediated eosinophil survival in these diseases. However,  
25 prolonged use of glucocorticoids can lead to side effects such as glaucoma, osteoporosis  
and growth retardation in the patients (see Hanania, N. A. et al., J. Allergy and Clin.  
Immunol., Vol. 96, 571-579 (1995) and Saha, M. T. et al., Acta Paediatrica, Vol. 86, #2,  
138-142 (1997)). It is therefore desirable to have an alternative means of treating

eosinophil related diseases without incurring these undesirable side effects.


Recently, the CCR-3 receptor was identified as a major chemokine receptor that eosinophils use for their response to eotaxin, RANTES and MCP-3. When transfected into a murine pre-beta. lymphoma line, CCR-3 bound eotaxin, RANTES and MCP-3 conferred chemotactic responses on these cells to eotaxin, RANTES and MCP-3 (see Ponath, P. D. et al., J. Exp. Med. 183, 2437-2448 (1996)). The CCR-3 receptor is expressed on the surface of eosinophils, T-cells (subtype Th-2), basophils and mast cells and is highly selective for eotaxin. Studies have shown that pretreatment of eosinophils with an anti-CCR-3 mAb completely inhibits eosinophil chemotaxis to eotaxin, RANTES and MCP-3 (see Heath, H. et al., J. Clin. Invest., Vol. 99, #2, 178-184 (1997)). Applicants' issued U.S. patents U.S. Patent Nos. 6,140,344 and 6,166,015 and published EP application EP903349, published March 24, 1999 disclose CCR-3 antagonists that inhibit eosinophilic recruitment by chemokine such as eotaxin.

Therefore, blocking the ability of the CCR-3 receptor to bind RANTES, MCP-3 and eotaxin and thereby preventing the recruitment of eosinophils should provide for the treatment of eosinophil-mediated inflammatory diseases.

The present invention concerns novel piperidine derivatives which are capable of inhibiting the binding of eotaxin to the CCR-3 receptor and thereby provide a means of combating eosinophil induced diseases, such as asthma.

20

In a first aspect, this invention provides a compound of Formula (I):



(I)

wherein:

25 R<sup>1</sup> is (C<sub>1</sub>-C<sub>2</sub>)alkylene;

R<sup>2</sup> is optionally substituted phenyl;

R<sup>3</sup> is hydrogen, alkyl, acyl, aryl, or arylalkyl;

- 3 -

ring A is a cycloalkyl, heterocyclyl, or optionally substituted phenyl;

L is  $-C(=O)-$ ,  $-C(=S)-$ ,  $-SO_2-$ ,  $-C(=O)N(R_a)-$ ,  $-C(=S)N(R_a)-$ ,  $-SO_2N(R_a)-$ ,  $-C(=O)O-$ ,  $-C(=S)O-$ ,  $-S(=O)_2O-$ ;

5 where  $R_a$  is hydrogen, alkyl, acyl, aryl, arylalkyl, alkoxy carbonyl, or benzyloxycarbonyl;

X is absent,  $-(CR'R'')O-$ ,  $-(CR'R'')S-$ ,  $-(CR'R'')NR_b-$  or alkylene;

where  $R'$  and  $R''$  are independently hydrogen or alkyl, and  $R_b$  is hydrogen or alkyl;

$R^4$  is aryl or heteroaryl; and


10  $R^5$  is hydrogen or alkyl;

provided that when  $R^1$  is  $-CH_2-$ ,  $R^2$  is phenyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, A is phenyl, L is  $-C(=O)NH-$  and X is absent, then  $R^4$  is not 2,5-difluorophenyl; or

prodrugs, individual isomers, racemic and non-racemic mixtures of isomers, and pharmaceutically acceptable salts thereof.

15 Also, within the compounds as defined above [they will be referred to in the following under (i)], preferred are the following compounds:

(ii) The compound of (i), which is a compound of Formula (II):




(II)

20 wherein  $R^1-R^5$ , A, L, and X are as defined in (i).

(iii) The compound of (i), which is a compound of Formula (III):


- 4 -



(III)

wherein R<sup>1</sup>-R<sup>5</sup>, A, L, and X are as defined in (i).

- (iv) The compound of any one of (i) to (iii) wherein R<sup>1</sup> is methylene.
- 5 (v) The compound of any one of (i) to (iii) wherein R<sup>2</sup> is 4-chlorophenyl or 3,4-dichlorophenyl.
- (vi) The compound of any one of (i) to (iii) wherein R<sup>3</sup> is hydrogen.
- (vii) The compound of any one of (i) to (iii) wherein L is -C(=O)-, -SO<sub>2</sub>-, -C(=O)N(R<sub>a</sub>)-, -C(=S)N(R<sub>a</sub>)-, or -C(=O)O-.
- 10 (viii) The compound of (vii) wherein L is -C(=O)-.
- (ix) The compound of (i), which is a compound of Formula (IV):




(IV)

wherein R<sup>3</sup>, R<sup>4</sup>, A, L, and X are as defined in (i).

- 15 (x) The compound of (i), which is a compound of Formula (VI):

- 5 -



(VI)

wherein X and R<sup>4</sup> are as defined in (i).

- (xi) The compound of (vii) wherein X is absent, methylene, 1,2-ethanediyl, 1,3-  
 5 propanediyl, or 1,4-butanediyl.
- (xii) The compound of (xi) wherein R<sup>4</sup> is 3,4-dichlorophenyl, 3,4,5-trimethoxyphenyl, 4-methanesulfonylphenyl, 3-methanesulfonylphenyl, 4-methoxynaphthalen-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, phenyl, 3-fluorophenyl, 4-ethylphenyl, 3-methoxyphenyl, 2,4-difluorophenyl, 3-trifluoromethylphenyl, 4-methylphenyl, 4-  
 10 fluorophenyl, 2-fluorophenyl, 4-carbamoylphenyl, 3-carbamoylphenyl, 4-acetylphenyl, 4-nitrophenyl, 2-methylphenyl, 2-chloro-4-fluorophenyl, 3,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 4-bromophenyl, 4-chloro-3-nitrophenyl, 2-nitrophenyl, 2-nitro-4-trifluoromethylphenyl, 4-chlorophenyl, 3-chlorophenyl, 2-chlorophenyl, 3-methylphenyl, 2-trifluoromethylphenyl, 2-  
 15 methoxyphenyl, 3-bromophenyl, 4-trifluoromethylphenyl, 3-trifluoromethylphenyl, 3,5-bis-trifluoromethylphenyl, 4-*tert*-butylphenyl, 4-ethoxyphenyl, 3-cyanophenyl, 4-cyanophenyl, 4-methoxyphenyl, 3-nitrophenyl, 3,5-dimethoxyphenyl, 4-iodophenyl, 4-isopropylphenyl, 3-methoxycarbonylphenyl, 3-acetylphenyl, 2-methylphenyl, indol-2-yl, 5-methoxyindol-2-yl, 5-chloroindol-2-yl, 2-methoxycarbonylphenyl, 3,5-dichlorophenyl,  
 20 1-naphthyl, 3-chloro-2-methylphenyl, 2,5-dimethylphenyl, 2-thienyl, 3-ethoxyphenyl, 3-isoquinolyl, 2-methylquinolin-6-yl, 3-methylaminophenyl, 3-quinolyl, 2-quinolyl, 5-hydroxynaphthalen-2-yl, 8-hydroxyquinolin-2-yl, 5,7-dimethyl-[1,8]naphthyridin-2-yl, 6-quinolyl, 3-(acetilamino)phenyl, or 2,3,4-trimethoxyphenyl.
- (xiii) The compound of (xii) wherein R<sup>4</sup> is 3,4,5-trimethoxyphenyl, 4-acetylphenyl, 3-  
 25 carbamoylphenyl, 4-carbamoylphenyl, 3-methanesulfonylphenyl or 4-methanesulfonylphenyl.

(xiv) The compound of (viii), wherein X is  $-\text{CH}_2\text{S}-$  and  $\text{R}^4$  is 5-(3,4-dimethoxyphenyl)-pyrimidin-2-yl, 5-(3,4-methylenedioxyphenyl)-pyrimidin-2-yl or 5-(4-methoxyphenyl)pyrimidin-2-yl, and a salt thereof.

(xv) The compound of (i), wherein ring A is cycloalkyl or heterocyclyl or substituted  
5 phenyl.

(xvi) The compound of (xv), wherein ring A is cycloalkyl or heterocyclyl.

(xvii) The compound of (i), wherein  $\text{R}^2$  is substituted phenyl.

In a second aspect, this invention provides pharmaceutical compositions containing a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically  
10 acceptable salt thereof, and a pharmaceutically acceptable excipient.

In a third aspect, this invention provides processes disclosed herein for preparing compounds of Formula (I).

In a forth aspect, this invention provides novel intermediates disclosed herein that are useful for preparing compounds of Formula (I).

15 In a fifth aspect, this invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof for use in medical therapy or diagnosis (e.g. for treating asthma).

In a sixth aspect, this invention provides the use of a compound of Formula (I) or a pharmaceutically acceptable salt thereof for the manufacture of a medicament useful for  
20 treating a disease in a mammal treatable by administration of a CCR-3 receptor antagonist (e.g. asthma).

Unless otherwise stated, the following terms used in the specification and claims have the meanings given below.

25 "Acyl" means a radical  $-\text{C}(\text{O})\text{R}$ , where R is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl wherein alkyl, cycloalkyl, cycloalkylalkyl, and phenylalkyl are as defined herein. Representative examples include, but are not limited to formyl, acetyl, cyclohexylcarbonyl, cyclohexylmethylcarbonyl, benzoyl, benzylcarbonyl.

"Acylalkyl" means a radical  $-\text{alkylene}-\text{C}(\text{O})\text{R}$  where R is hydrogen, alkyl, haloalkyl,  
30 cycloalkyl, cycloalkyl-alkyl, optionally substituted phenyl, benzyl, hydroxy, alkoxy, amino,

monoalkylamino or dialkylamino. Representative examples include methylcarbonylmethyl, 2-(ethoxycarbonyl)ethyl, 2-(methoxycarbonyl)ethyl, 2-carboxyethyl.

"Acylamino" means a radical  $-NR'C(O)R$ , where R' is hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl wherein alkyl, cycloalkyl, cycloalkylalkyl, and phenylalkyl are as defined herein. Representative examples include, but are not limited to formylamino, acetylamino, cyclohexylcarbonylamino, cyclohexylmethyl-carbonylamino, benzoylamino, benzylcarbonylamino.

"Alkoxy" means a radical  $-OR$  where R is an alkyl as defined herein e.g., methoxy, ethoxy, propoxy, butoxy.

10 "Alkoxycarbonyl" means a radical  $-C(O)-R$  where R is alkoxy as defined herein.

"Alkenyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenyl, propenyl.

15 "Alkyl" means a linear saturated monovalent hydrocarbon radical of one to six carbon atoms or a branched saturated monovalent hydrocarbon radical of three to six carbon atoms, e.g., methyl, ethyl, propyl, 2-propyl, *n*-butyl, *iso*-butyl, *tert*-butyl, pentyl.

"Alkylamino" or "Monoalkylamino" means a radical  $-NHR$  where R represents an alkyl, cycloalkyl or cycloalkyl-alkyl group as defined herein. Representative examples include, but are not limited to methylamino, ethylamino, isopropylamino, cyclohexylamino.

"Alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., methylene, ethylene, 2,2-dimethylethylene, propylene, 2-methylpropylene, butylene, pentylene.

25 "Alkynyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one triple bond, e.g., ethynyl, propynyl.

"Alkylsulfonyl" means a radical  $-S(O)_2R$  where R is an alkyl, cycloalkyl or cycloalkyl-alkyl group as defined herein, e.g., methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl, cyclohexylsulfonyl.



"Alkylsulfinyl" means a radical  $-S(O)R$  where R is an alkyl, cycloalkyl or cycloalkyl-alkyl group as defined herein e.g., methylsulfinyl, ethylsulfinyl, propylsulfinyl, butylsulfinyl, cyclohexylsulfinyl.

"Alkylthio " means a radical  $-SR$  where R is an alkyl as defined above e.g., methylthio, ethylthio, propylthio, butylthio.

"Aryl" means a monocyclic or bicyclic aromatic hydrocarbon radical of preferably 6 to 10 ring atoms which is optionally substituted with one or more substituents, preferably one, two or three, substituents preferably selected from the group consisting of alkyl, haloalkyl, hydroxyalkyl, heteroalkyl, acyl, acylamino, amino, alkylamino, dialkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl,  $-SO_2NR'R''$  (where R' and R'' are independently hydrogen or alkyl), alkoxy, haloalkoxy, alkoxycarbonyl, carbamoyl, hydroxy, halo, nitro, cyano, mercapto, methylenedioxy or ethylenedioxy. More specifically the term aryl includes, but is not limited to, phenyl, chlorophenyl, fluorophenyl, methoxyphenyl, 1-naphthyl, 2-naphthyl, and the derivatives thereof.

"Arylene" means a divalent aryl group as defined above.

"Arylalkyl" refers to an alkyl radical as defined herein in which one of the hydrogen atoms of the alkyl group is replaced with an aryl group. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl.

"Aryloxy" means a radical  $-O-R$  where R is an aryl group as defined herein.

"Carbamoyl" means the radical  $-C(=O)NH_2$ .

"Cycloalkyl" refers to a saturated monovalent cyclic hydrocarbon radical of three to seven ring carbons e.g., cyclopropyl, cyclobutyl, cyclohexyl, 4-methylcyclohexyl.

"Cycloalkyl-alkyl" means a radical  $-R^xR^y$  where  $R^x$  is an alkylene group and  $R^y$  is cycloalkyl group as defined herein, e.g., cyclohexylmethyl.

"Dialkylamino" means a radical  $-NRR'$  where R and R' independently represent an alkyl, cycloalkyl, or cycloalkylalkyl group as defined herein. Representative examples include, but are not limited to dimethylamino, methylethylamino, di(1-methylethyl)amino, (cyclohexyl)(methyl)amino, (cyclohexyl)(ethyl)amino, (cyclohexyl)(propyl)amino, (cyclohexylmethyl)(methyl)amino, (cyclohexylmethyl)(ethyl)amino.

"Halo" means fluoro, chloro, bromo, or iodo, preferably fluoro and chloro.

"Haloalkyl" means alkyl substituted with one or more same or different halo atoms, e.g.,  $-\text{CH}_2\text{Cl}$ ,  $-\text{CF}_3$ ,  $-\text{CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CCl}_3$ .

"Heteroaryl" means a monocyclic or bicyclic radical of 5 to 12 ring atoms having at least one aromatic ring containing one, two, or three ring heteroatoms selected from N, O, or S, the remaining ring atoms being C, with the understanding that the attachment point of the heteroaryl radical will be on an aromatic ring. The heteroaryl ring is optionally substituted independently with one or more substituents, preferably one or two substituents, selected from alkyl, haloalkyl, hydroxyalkyl, heteroalkyl, acyl, acylamino, amino, alkylamino, dialkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl,  $-\text{SO}_2\text{NR}'\text{R}''$  (where R' and R'' are independently hydrogen or alkyl), alkoxy, haloalkoxy, alkoxycarbonyl, carbamoyl, hydroxy, halo, nitro, cyano, mercapto, methylenedioxy, ethylenedioxy or optionally substituted phenyl. More specifically the term heteroaryl includes, but is not limited to, pyridyl, furanyl, thienyl, thiazolyl, isothiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrrolyl, pyrazolyl, pyrimidinyl, 5-(3,4-dimethoxyphenyl)-pyrimidin-2-yl, 5-(4-methoxyphenyl)-pyrimidin-2-yl, 5-(3,4-methylenedioxyphenyl)-pyrimidin-2-yl, benzofuranyl, tetrahydrobenzofuranyl, isobenzofuranyl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl, indolyl, isoindolyl, benzoxazolyl, quinolyl, tetrahydroquinolyl, isoquinolyl, benzimidazolyl, benzisoxazolyl or benzothienyl and derivatives thereof.

"Heteroarylene" means a divalent heteroaryl group as defined above.

"Heteroarylalkyl" means an alkyl radical as defined herein in which one of the hydrogen atoms of the alkyl group is replaced with a heteroaryl group.

"Heteroalkyl" means an alkyl radical as defined herein wherein one, two or three hydrogen atoms have been replaced with a substituent independently selected from the group consisting of  $-\text{OR}^a$ ,  $-\text{NR}^b\text{R}^c$ , and  $-\text{S}(\text{O})_n\text{R}^d$  (where n is an integer from 0 to 2), with the understanding that the point of attachment of the heteroalkyl radical is through a carbon atom, wherein  $\text{R}^a$  is hydrogen, acyl, alkyl, cycloalkyl, or cycloalkylalkyl;  $\text{R}^b$  and  $\text{R}^c$  are independently of each other hydrogen, acyl, alkyl, cycloalkyl, or cycloalkylalkyl; when n is 0,  $\text{R}^d$  is hydrogen, alkyl, cycloalkyl, or cycloalkylalkyl, and when n is 1 or 2,  $\text{R}^d$  is alkyl, cycloalkyl, cycloalkylalkyl, amino, acylamino, monoalkylamino, or dialkylamino; Representative examples include, but are not limited to, 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxy-1-hydroxymethylethyl, 2,3-dihydroxypropyl, 1-hydroxymethylethyl, 3-hydroxybutyl, 2,3-dihydroxybutyl, 2-hydroxy-1-methylpropyl, 2-aminoethyl, 3-aminopropyl, 2-methylsulfonylethyl, aminosulfonylmethyl, aminosulfonylethyl,

aminosulfonylpropyl, methylaminosulfonylmethyl, methylaminosulfonylethyl, methylaminosulfonylpropyl.

"Heterocyclyl" means a saturated or unsaturated non-aromatic cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from  $\text{NR}^x$  {wherein  
 5 each  $\text{R}^x$  is independently hydrogen, alkyl, acyl, alkylsulfonyl, aminosulfonyl, (alkylamino)sulfonyl, (dialkylamino)sulfonyl, carbamoyl, (alkylamino)carbonyl, (dialkylamino)carbonyl, (carbamoyl)alkyl, (alkylamino)carbonylalkyl, or dialkylaminocarbonylalkyl}, O, or  $\text{S}(\text{O})_n$  (where n is an integer from 0 to 2), the remaining ring atoms being C. The heterocyclyl ring may be optionally substituted independently  
 10 with one, two, or three substituents selected from alkyl, haloalkyl, heteroalkyl, halo, nitro, cyanoalkyl, hydroxy, alkoxy, amino, monoalkylamino, dialkylamino, arylalkyl,  $-(\text{X})_n-\text{C}(\text{O})\text{R}$  (where X is O or  $\text{NR}'$ , n is 0 or 1, R is hydrogen, alkyl, haloalkyl, hydroxy, alkoxy, amino, monoalkylamino, dialkylamino or optionally substituted phenyl, and  $\text{R}'$  is hydrogen or alkyl),  $-\text{alkylene}-\text{C}(\text{O})\text{R}$  (where R is hydrogen, alkyl, haloalkyl, hydroxy, alkoxy, amino, monoalkylamino, dialkylamino or optionally substituted phenyl) or  
 15  $-\text{S}(\text{O})_n\text{R}^d$  (where n is an integer from 0 to 2, and  $\text{R}^d$  is hydrogen (provided that n is 0), alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, amino, monoalkylamino, dialkylamino, or hydroxyalkyl). More specifically the term heterocyclyl includes, but is not limited to, tetrahydropyranyl, piperidino, N-methylpiperidin-3-yl, piperazino, N-methylpyrrolidin-3-yl, 3-pyrrolidino, morpholino, thiomorpholino, thiomorpholino-1-oxide,  
 20 thiomorpholino-1,1-dioxide, tetrahydrothiophenyl-S,S-dioxide, pyrrolinyl, imidazoliny, and the derivatives thereof.

"Hydroxyalkyl" means an alkyl radical as defined herein, substituted with one or more, preferably one, two or three hydroxy groups, provided that the same carbon atom  
 25 does not carry more than one hydroxy group. Representative examples include, but are not limited to, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-(hydroxymethyl)-2-methylpropyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl, 2,3-dihydroxypropyl, 2-hydroxy-1-hydroxymethylethyl, 2,3-dihydroxybutyl, 3,4-dihydroxybutyl and 2-(hydroxymethyl)-3-hydroxypropyl, preferably 2-hydroxyethyl, 2,3-dihydroxypropyl and  
 30 1-(hydroxymethyl)-2-hydroxyethyl. Accordingly, as used herein, the term "hydroxyalkyl" is used to define a subset of heteroalkyl groups.

"Leaving group" has the meaning conventionally associated with it in synthetic organic chemistry, i.e., an atom or a group capable of being displaced by a nucleophile and includes halo (such as chloro, bromo, and iodo), alkanesulfonyloxy, arenesulfonyloxy,

alkylcarbonyloxy (e.g., acetoxy), arylcarbonyloxy, mesyloxy, tosyloxy, trifluoromethanesulfonyloxy, aryloxy (e.g., 2,4-dinitrophenoxy), methoxy, N,O-dimethylhydroxylamino.

"Optionally substituted phenyl" means a phenyl group which is optionally substituted with one or more substituents, preferably one, two or three, substituents preferably selected from the group consisting of alkyl, haloalkyl, hydroxyalkyl, heteroalkyl, acyl, acylamino, amino, alkylamino, dialkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, -SO<sub>2</sub>NR'R'' (where R' and R'' are independently hydrogen or alkyl), alkoxy, haloalkoxy, alkoxycarbonyl, carbamoyl, hydroxy, halo, nitro, cyano, mercapto, methylenedioxy or ethylenedioxy. More specifically the term includes, but is not limited to, phenyl, chlorophenyl, fluorophenyl, bromophenyl, methylphenyl, ethylphenyl, methoxyphenyl, cyanophenyl, 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-methylphenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl and the derivatives thereof.

"Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "aryl group optionally mono- or di-substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the aryl group is mono- or disubstituted with an alkyl group and situations where the aryl group is not substituted with the alkyl group.

"Pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes excipient that is acceptable for veterinary use as well as human pharmaceutical use. A "pharmaceutically acceptable excipient" as used in the specification and claims includes both one and more than one such excipient.

"Pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid,

succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine.

"Phenylalkyl" refers to an alkyl radical as defined herein in which one of the hydrogen atoms of the alkyl radical has been replaced by an optionally substituted phenyl.

"Protecting group" refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity. Examples of protecting groups can be found in T.W. Green and P.G. Futs, Protective Groups in Organic Chemistry, (Wiley, 2<sup>nd</sup> ed. 1991) and Harrison and Harrison et al., Compendium of Synthetic Organic Methods, Vols. 1-8 (John Wiley and Sons, 1971-1996). Representative amino protecting groups include, formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (CBZ), *tert*-butoxycarbonyl (Boc), trimethyl silyl (TMS), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (Fmoc), nitro-veratryloxycarbonyl (NVOC). Representative hydroxy protecting groups include those where the hydroxy group is either acylated or alkylated such as benzyl, and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers.

"Treating" or "treatment" of a disease includes: (1) preventing the disease, i.e., causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease; (2) inhibiting the disease, i.e., arresting or reducing the development of the disease or its clinical symptoms; or (3) relieving the disease, i.e., causing regression of the disease or its clinical symptoms.

"A therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for

the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

"Prodrugs" means any compound which releases an active parent drug according to Formula I *in vivo* when such a prodrug is administered to a mammalian subject. Prodrugs of a compound of Formula I are prepared by modifying functional groups present in the compound of Formula I in such a way that the modifications may be cleaved *in vivo* to release the parent compound. Prodrugs include compounds of Formula I wherein a hydroxy, amino, or sulfhydryl group in a compound of Formula I is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to esters (e.g., acetate, formate, and benzoate derivatives), carbamates (e.g., N,N-dimethylaminocarbonyl) of hydroxy functional groups in compounds of Formula I.






Compounds that have the same molecular Formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers." Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers". When a compound has an asymmetric center, for example, if a carbon atom is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the R- and S-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture".






The compounds of this invention may possess one or more asymmetric centers; such compounds can therefore be produced as individual (R)- or (S)-stereoisomers or as mixtures thereof. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition J. March, John Wiley and Sons, New York, 1992). The nomenclature used in this application is generally based on the IUPAC recommendations. For example, a compound of Formula (I) wherein R<sub>1</sub> is methylene; R<sub>2</sub> is 4-chlorophenyl; L is C(=O)NH; X is absent; A is

cyclopentyl; R<sub>3</sub> is hydrogen; and R<sub>4</sub> is 2-quinolyl (Compound 1 in Table 1), is named (±)-*trans*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]cyclopentyl}-3-quinolin-2-yl-urea.

Representative compounds of Formula (I) are shown in the following table.

Table I

Compound	Structure	M.P. (°C)
1		
2		
3		
4		
5		

Compound	Structure	M.P. (°C)
6		
7		
8		
9		
10		

While the broadest definition of this invention is described before, certain compounds of Formula (I) are preferred.

A preferred compound of the invention is a compound of Formula (I) wherein R<sup>1</sup> is  
 5 methylene.

One aspect of the invention relates to compounds of Formula (I) where ring A is cycloalkyl, heterocyclyl or substituted phenyl. Another preferred compound of the invention are compounds of Formula (I) wherein ring A is cyclopentyl. Compounds where ring A is cyclopentyl are bind unexpectedly potently to the CCR-3 receptor. Other



preferred compounds of the invention are compounds of Formula (I) wherein ring A is heterocyclyl (particularly tetrahydropyranyl, S,S-dioxo-tetrahydrothiophenyl, tetrahydrothiophenyl or pyrrolidinyl) or compounds of Formula (I) wherein ring A is substituted phenyl.

5 A preferred compound of the invention is a compound of Formula (I) wherein R<sup>2</sup> is phenyl ring substituted with one, or two substituents selected from alkyl, alkoxy, haloalkyl, halo, cyano or nitro; preferably methyl, ethyl, methoxy, trifluoromethyl, chloro, fluoro or bromo; most preferably 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-methylphenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl. Particularly preferred are 4-chlorophenyl or  
10 3,4-dichlorophenyl.

A preferred compound of the invention is a compound of Formula (I) wherein R<sup>3</sup> is hydrogen or methyl, preferably hydrogen.

A preferred compound of the invention is a compound of Formula (I) wherein L is -  
15 C(=O)-, -SO<sub>2</sub>-, -C(=O)N(R<sub>a</sub>)-, -C(=S)N(R<sub>a</sub>)-, or -C(=O)O-. More preferred are compounds where L is -C(=O)-, -C(=O)N(R<sub>a</sub>)-, most preferably -C(=O)N(R<sub>a</sub>)-. In the preceding R<sub>a</sub> is preferably hydrogen or methyl, most preferably hydrogen.


A preferred compound of the invention is a compound of Formula (I) wherein X is absent, methylene, 1,2-ethanediyl, 1,3-propanediyl, or 1,4-butanediyl.

20 A preferred compound of the invention is a compound of Formula (I) wherein R<sup>4</sup> is optionally substituted phenyl, optionally substituted heteroaryl wherein the heteroaryl ring is indolyl, thienyl, quinoliny, substituted pyrimidin-2-yl, e.g. (5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, 5-(3,4-methylenedioxy)-pyrimidin-2-yl or 5-(4-methoxyphenyl)pyrimidin-2-yl) or 1,8-naphthyridinyl. Preferably R<sup>4</sup> is selected from 3,4-dichlorophenyl, 3,4,5-trimethoxyphenyl, 4-methanesulfonyl-phenyl, 3-methanesulfonylphenyl, 4-methoxynaphthalen-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, phenyl, 3-fluorophenyl, 4-ethylphenyl, 3-methoxyphenyl, 2,4-difluorophenyl, 3-trifluoromethylphenyl, 4-methylphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-carboxamidophenyl, 4-acetylphenyl, 4-nitrophenyl, 2-methylphenyl, 2-chloro-4-fluorophenyl, 3,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 2,3-dichlorophenyl,  
25 2,4-dichlorophenyl, 4-bromophenyl, 4-chloro-3-nitrophenyl, 2-nitrophenyl, 2-nitro-4-trifluoromethylphenyl, 4-chlorophenyl, 3-chlorophenyl, 2-chlorophenyl, 3-methylphenyl, 2-trifluoromethylphenyl, 2-methoxyphenyl, 3-bromophenyl, 4-trifluoromethylphenyl, 3-

trifluoromethylphenyl, 3,5-bis-trifluoromethylphenyl, 4-*tert*-butylphenyl, 4-ethoxyphenyl, 3-cyanophenyl, 4-cyanophenyl, 4-methoxyphenyl, 3-nitrophenyl, 3,5-dimethoxyphenyl, 4-iodophenyl, 4-isopropylphenyl, 3-methoxycarbonylphenyl, 3-acetylphenyl, 2-methylphenyl, indol-2-yl, 5-methoxyindol-2-yl, 5-chloroindol-2-yl, 2-methoxycarbonylphenyl, 3,5-dichlorophenyl, 1-naphthyl, 3-chloro-2-methylphenyl, 2,5-dimethylphenyl, 2-thienyl, 3-ethoxyphenyl, 3-isoquinolyl, 2-methylquinolin-6-yl, 3-methylaminophenyl, 3-quinolyl, 2-quinolyl, 5-hydroxynaphthalen-2-yl, 8-hydroxyquinolin-2-yl, 5,7-dimethyl-[1,8]naphthyridin-2-yl, 6-quinolyl, 3-(acetylamino)phenyl, or 2,3,4-trimethoxyphenyl. Particularly preferred are R<sup>4</sup> being trimethoxyphenyl, e.g 3,4,5-trimethoxyphenyl, 4-acetyl-phenyl, 4-carboxamido-phenyl and 4-methanesulfonyl-phenyl.


Also preferred are compounds where X is -CH<sub>2</sub>S-, -CH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>- and R<sup>4</sup> is heteroaryl, preferably optionally substituted pyrimidinyl, pyrazolyl or thienyl. Particularly preferred are compounds where X is -CH<sub>2</sub>S- and R<sup>4</sup> is 5-(3,4-dimethoxyphenyl)-pyrimidin-2-yl, 5-(3,4-methylenedioxyphenyl)-pyrimidin-2-yl, 5-(4-methoxyphenyl)pyrimidin-2-yl

A specific compound of Formula (I) is a compound of Formula (II):




wherein R<sup>1</sup>-R<sup>5</sup>, A, L, and X have any of the values described herein.

A specific compound of Formula (I) is a compound of Formula (III):




wherein R<sup>1</sup>-R<sup>5</sup>, A, L, and X have any of the values described herein.

A specific compound of Formula (I) is a compound of formula (IV):



wherein  $R^3$ ,  $R^4$ , A, L, and X have any of the values described herein.


A specific compound of Formula (I) is a compound of formula (V):



5

wherein X and  $R^4$  have any of the values defined herein.

A specific compound of Formula (I) is a compound of formula (VI):




wherein X and  $R^4$  have any of the values defined herein.

10


A specific compound of Formula (I) is a compound of formula (VII):

- 19 -




wherein X and R<sup>4</sup> have any of the values defined herein.

A specific compound of Formula (I) is a compound of formula (VIII):



- 5 wherein X and R<sup>4</sup> have any of the values defined herein; and R<sup>x</sup> is hydrogen, alkyl, acyl, alkylsulfonyl, aminosulfonyl, (alkylamino)sulfonyl, (dialkylamino)sulfonyl, carbamoyl, (alkylamino)carbonyl, (dialkylamino)carbonyl, (carbamoyl)alkyl, (alkylamino)carbonylalkyl, or dialkylaminocarbonylalkyl.

A specific compound of Formula (I) is a compound of formula (IX):




10

wherein X and R<sup>4</sup> have any of the values defined herein.

A specific compound of Formula (I) is a compound of formula (X):

- 20 -



wherein X and R<sup>4</sup> have any of the values defined herein.

A particularly preferred compound of the invention is:

*trans*-1-{2-[4-(4-chlorobenzyl)-piperidin-1-yl]-cyclopentyl}-3-(3,4,5-trimethoxy-  
5 phenyl)urea hydrochloride;

*trans*-1-{4-[4-(4-chlorobenzyl)-piperidin-1-yl]-tetrahydro-furan-3-yl}-3-(3,4,5-trimethoxy-phenyl)urea;

(±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxy-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea;

10 (±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxymethyl-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea; and

(±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-methoxymethyl-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea

or prodrugs, individual isomers, racemic and non-racemic mixtures of isomers, and  
15 pharmaceutically acceptable salts thereof.

The compounds of the invention are CCR-3 receptor antagonists and inhibit eosinophil recruitment by CCR-3 chemokines such as RANTES, eotaxin, MCP-2, MCP-3 and MCP-4. Compounds of this invention and compositions containing them are useful in the treatment of eosinophil-induced diseases such as inflammatory or allergic diseases  
20 and including respiratory allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, hypersensitivity pneumonitis, eosinophilic pneumonias (e.g., chronic eosinophilic pneumonia); inflammatory bowel diseases (e.g., Crohn's disease and ulcerative colitis); and psoriasis and inflammatory dermatoses such as dermatitis and eczema.

The CCR-3 antagonistic activity of the compounds of this invention can be measured by *in vitro* assays such as ligand binding and chemotaxis assays as described in more detail in Examples 45, 46, and 47. *In vivo* activity was assayed in the Ovalbumin induced Asthma in Balb/c Mice Model as described in more detail in Example 48.

5

In general, the compounds of this invention can be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. The actual amount of the compound of this invention, i.e., the active ingredient, will depend upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, and other factors.

Therapeutically effective amounts of compounds of Formula (I) may range from approximately 0.01-20 mg per kilogram body weight of the recipient per day; preferably about 0.1-10 mg/kg/day. Thus, for administration to a 70 kg person, the dosage range would most preferably be about 7 mg to 0.7 g per day.

15

In general, compounds of this invention will be administered as pharmaceutical compositions by any one of the following routes: oral, transdermal, inhalation (e.g., intranasal or oral inhalation) or parenteral (e.g., intramuscular, intravenous or subcutaneous) administration. A preferred manner of administration is oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction. Compositions can take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, liposomes, elixirs, or any other appropriate compositions. Another preferred manner for administering compounds of this invention is inhalation. This is an effective means for delivering a therapeutic agent directly to the respiratory tract for the treatment of diseases such as asthma and other similar or related respiratory tract disorders (see U.S. Pat. No. 5,607,915).

20

The choice of formulation depends on various factors such as the mode of drug administration and the bioavailability of the drug substance. For delivery via inhalation the compound can be formulated as liquid solutions or suspensions, aerosol propellants or dry powder and loaded into a suitable dispenser for administration. There are three types of pharmaceutical inhalation devices--nebulizer inhalers, metered-dose inhalers (MDI) and dry powder inhalers (DPI). Nebulizer devices produce a stream of high velocity air that causes the therapeutic agents (which has been formulated in a liquid form) to spray as a mist which is carried into the patient's respiratory tract. MDI's typically have the formulation packaged with a compressed gas. Upon actuation, the device discharges a

35

measured amount of therapeutic agent by compressed gas, thus affording a reliable method of administering a set amount of agent. DPI's administer therapeutic agents in the form of a free flowing powder that can be dispersed in the patient's inspiratory air-stream during breathing by the device. In order to achieve a free flowing powder, the therapeutic agent is formulated with an excipient, such as lactose. A measured amount of the therapeutic is stored in a capsule form and is dispensed to the patient with each actuation. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability can be increased by increasing the surface area i.e., decreasing particle size. For example, U.S. Pat. No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a crosslinked matrix of macromolecules. U.S. Pat. No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

The compositions are comprised of in general, a compound of Formula (I) in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the compound of Formula (I). Such excipient may be any solid, liquid, semi-solid or, in the case of an aerosol composition, gaseous excipient that is generally available to one of skill in the art.

Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk. Liquid and semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

Compressed gases may be used to disperse a compound of this invention in aerosol form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.

For liposomal formulations of the drug for parenteral or oral delivery the drug and the lipids are dissolved in a suitable organic solvent e.g. tert-butanol, cyclohexane (1% ethanol). The solution is lyophilized and the lipid mixture is suspended in an aqueous buffer and allowed to form a liposome. If necessary, the liposome size can be reduced by

sonification. (see, Frank Szoka, Jr. and Demetrios Papahadjopoulos, "Comparative Properties and Methods of Preparation of Lipid Vesicles (Liposomes)", *Ann. Rev. Biophys. Bioeng.*, 9:467-508 (1980), and D. D. Lasic, "Novel Applications of Liposomes", *Trends in Biotech.*, 16:467-608, (1998)).

- 5           Other suitable pharmaceutical excipients and their formulations are described in Remington's Pharmaceutical Sciences, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).

10           The level of the compound in a formulation can vary within the full range employed by those skilled in the art. Typically, the formulation will contain, on a weight percent (wt %) basis, from about 0.01-99.99 wt % of a compound of Formula (I) based on the total formulation, with the balance being one or more suitable pharmaceutical excipients. Preferably, the compound is present at a level of about 1-80 wt %. Representative pharmaceutical formulations containing a compound of Formula (I) are described in Example 44.

15

          The compounds of the present invention can be prepared in a number of ways known to one skilled in the art. Preferred methods include, but are not limited to, the general synthetic procedures described below.

20           The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, Wis., USA), Bachem (Torrance, Calif., USA), Enika Chemie, or Sigma (St. Louis, Mo., USA), Maybridge (Dist: Ryan Scientific, P.O. Box 6496, Columbia, S.C. 92960), Bionet Research Ltd., (Cornwall PL32 9QZ, UK), Menai Organics Ltd., (Gwynedd, N. Wales, UK), Butt Park Ltd., (Dist. Interchim, Montlucon Cedex, France) or are prepared by methods known to those skilled in the art following procedures set forth in references such as Fieser and Fieser's Reagents for Organic Synthesis, Volumes 1-17 (John Wiley and Sons, 1991); Rodd's Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989), Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991), March's Advanced Organic Chemistry, (John Wiley and Sons, 1992), and Larock's  
25           Comprehensive Organic Transformations (VCH Publishers Inc., 1989). These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure.


          The starting materials and the intermediates of the reaction may be isolated and



purified if desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography. Such materials may be characterized using conventional means, including physical constants and spectral data.

## 5 Synthesis of Compounds of Formula (I)


Compounds of Formula (I) are generally prepared from the precursor amine of Formula (Ia) as shown below.




10 Preparation of compounds of Formula (Ia) and their conversion to compounds of Formula I is illustrated in the following Schemes 1-8.

Schemes 1-5 show methods of preparing compounds of Formula Ia having different rings A. Specific exemplification is provided for R<sup>1</sup>-R<sup>2</sup> being 4-chlorobenzyl in Preparations 1-6. Preparation of analogous compounds where R<sup>1</sup> and R<sup>2</sup> vary within the  
 15 full scope of this invention may be readily prepared by one of skill in the art in light of this specification and incorporated references.


### Scheme 1. Synthesis of Cyclobutylamines – Ring A = Phenyl.



## Scheme 2. Synthesis of Cis Diamines - Ring A = Cyclopentyl and Cyclohexyl.



## 5 Scheme 3. Synthesis of Trans Diamines - Ring A = Cycloalkyl, Tetrahydrofuranyl, Pyrrolidinyl or Tetrahydrothiophenyl.


10 General Procedure A: (Amine Alkylation with Epoxides)

A 0.5-1.5 M solution of the amine,  $R_2NH$  (1 equiv), and the specified epoxide, 3a (1.1-10 equiv) in EtOH is stirred at 80-95 °C for 2-4.5 days, allowed to cool to room temperature, and concentrated. The crude amino alcohol is purified by chromatography or recrystallization.


15 General Procedure B: (Amine Formation Using Methanesulfonyl Chloride and Ammonium Hydroxide)

A 0.2-0.3 M solution of the amino alcohol (1 equiv) in  $CH_2Cl_2$  at 0 °C is treated successively with  $Et_3N$  (2 equiv) and  $MeSO_2Cl$  (2 equiv), stirred at 0 °C for 1-2 hours, and partitioned between  $CH_2Cl_2$  and 10-15%  $NH_4OH$ . The aqueous phase is extracted with  $CH_2Cl_2$  and the extracts are dried and concentrated. A 0.13M solution of the residue in 2.5:1 dioxane:28-30 wt %  $NH_4OH$  is stirred at 70-80 °C for 2.5-18 hours, allowed to cool to room temperature, and concentrated. The residue is partitioned between EtOAc and 1 N NaOH, the aqueous phase is extracted with EtOAc, and the extracts are washed with brine, dried and concentrated. The crude product is purified by chromatography or used without further purification.

## Scheme 4. Synthesis of Sulfone – Ring A = Sulfolane (IS THIS CORRECT??).




## 5 Scheme 5. Synthesis of Aniline – Ring A = Phenyl.




Schemes 6 and 7 show preparation of compounds of Formula Ia where ring A is substituted. Scheme 6 shows preparation of compounds of Formula Ia with a substituted cyclopentyl ring A. Scheme 7 shows preparation of compounds of Formula Ia with a substituted pyrrolidine ring A by treatment of the unsubstituted pyrrolidine 7a (R=H) with the appropriate reagent to produce the substituted pyrrolidine 7b.

## 15 Scheme 6. Synthesis of Cycloalkyl Derivatives.




## Scheme 7. Synthesis of Pyrrolidine Derivatives.



5 Schemes 8 and 9 show methods of converting compounds of Formula (Ia) to compounds of Formula (I) where L and A are varied.

## Scheme 8. Conversion of Primary Amines to Ureas and Benzamides.

10 Compounds of Formula (I) where L is  $-\text{C}(=\text{O})\text{NR}_a$  and X is absent are made as shown below in Scheme 8 (exemplified with  $\text{R}^4$  being 3,4,5-trimethoxyphenyl) and General Procedures C and D. Compounds of Formula (I) where L is  $-\text{C}(=\text{O})-$  and X is absent are made as shown below in Scheme 8 (exemplified with  $\text{R}^4$  being 3,4,5-trimethoxyphenyl) and General Procedures E and F.



15

General Procedure C: (Urea Formation Using Isocyanates)

A 0.1-0.6 M solution of the amine (1 equiv) in  $\text{CH}_2\text{Cl}_2$  or  $\text{CH}_2\text{Cl}_2$  and DMF at 0-20 °C is treated with the specified isocyanate (1.1-2 equiv), stirred for 0.5-1.5 hours, and partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase is extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts are dried and concentrated. The crude urea is purified by column chromatography or preparative TLC or used in the next step without further purification.

20

General Procedure D: (Urea Formation Using Isocyanates followed by salt formation)

A 0.1-0.6 M solution of the amine (1 equiv) in  $\text{CH}_2\text{Cl}_2$  or  $\text{CH}_2\text{Cl}_2$  and DMF at 0-20 °C is treated with the specified isocyanate (1.1-2 equiv), stirred for 0.5-1.5 hours, and partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase is extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts are dried and concentrated. The crude urea is purified by column chromatography or preparative TLC or used in the next step without further purification. A solution of the free base in  $\text{CH}_2\text{Cl}_2$  is treated with 1 N HCl in  $\text{Et}_2\text{O}$  and concentrated to give the hydrochloride salt.

10

General Procedure E: (Amide Formation Using 1-Hydroxybenzotriazole and 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide Hydrochloride)

A 0.1-0.4 M solution of the amine (1 equiv) and the specified carboxylic acid (1.2-1.5 equiv) in  $\text{CH}_2\text{Cl}_2$  at 0 °C is treated successively with 1-hydroxybenzotriazole hydrate (HOBt) (0.2-0.5 equiv) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (DEC) (1.3-2 equiv), stirred at 0-20 °C for 2-72 hours, and partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase is extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts are dried and concentrated. The crude amide is purified by column chromatography and/or preparative TLC.


20

General Procedure F: (Amide Formation Using 1-Hydroxybenzotriazole and 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide Hydrochloride followed by salt formation)

A 0.1-0.4 M solution of the amine (1 equiv) and the specified carboxylic acid (1.2-1.5 equiv) in  $\text{CH}_2\text{Cl}_2$  at 0 °C is treated successively with 1-hydroxybenzotriazole hydrate (HOBt) (0.2-0.5 equiv) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (DEC) (1.3-2 equiv), stirred at 0-20 °C for 2-72 hours, and partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase is extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts are dried and concentrated. The crude amide is purified by column chromatography and/or preparative TLC. A solution of the free base in  $\text{CH}_2\text{Cl}_2$  is treated with 1 N HCl in  $\text{Et}_2\text{O}$  and concentrated to provide the hydrochloride salt.

30

Scheme 9 and following procedures G-O describe the various methods used to convert compounds of Formula Ia to compounds of Formula I where L is varied.



#### General Procedure G (Parallel Synthesis of Sulfonamides)

A mixture of the requisite amine Ia (1 equiv), the appropriate sulfonyl chloride (1.5 equiv), and Amberlite IRA67 (2 equiv) in  $\text{CH}_2\text{Cl}_2$  (2 ml) was rotated overnight. The mixture was treated with PS-trisamine (1.2 equiv) (Argonaut Technologies Inc., San Carlos, CA, USA) and rotated overnight. The solid was collected by filtration and washed with  $\text{CH}_2\text{Cl}_2$ , MeOH, and  $\text{CH}_2\text{Cl}_2$ . The filtrate was concentrated to give the product.

#### General Procedure H (Parallel Synthesis of Amides from Acid Chlorides)

A mixture of the requisite amine Ia (1 equiv), the appropriate acid chloride (1.5 equiv), and Amberlite IRA67 (2 equiv) in  $\text{CH}_2\text{Cl}_2$  (2 ml) was rotated overnight. The mixture was treated with PS-trisamine (1.2 equiv) and MP-carbonate (2 equiv) (Argonaut Technologies, San Carlos, CA) and rotated overnight. The solid was collected by filtration and washed with  $\text{CH}_2\text{Cl}_2$ , MeOH, and  $\text{CH}_2\text{Cl}_2$ . The filtrate was concentrated to give the product.

General Procedure I (Parallel Synthesis of Amides from Carboxylic Acids)

A mixture of the requisite amine Ia (1 equiv), the appropriate carboxylic acid (1.5 equiv), and PS-carobodiimide (2 equiv) (Argonaut Technologies Inc., San Carlos, CA, USA) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was rotated overnight. The mixture was treated with  
5 MP-carbonate (2 equiv) and rotated overnight. The solid was collected by filtration and washed with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was concentrated to give the product.

General Procedure J (Parallel Synthesis of Ureas from Isocyanates and Purification by Parallel Chromatography)

10 A mixture of the requisite amine Ia (1 equiv) and the appropriate isocyanate (1.2 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was stirred overnight. The mixture was concentrated to give the crude product, which was purified by parallel chromatography using a step gradient (2.5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>, 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>).

General Procedure K (Parallel Synthesis of Ureas from Isocyanates and Purification by Catch and Release Scavenger)

15 A mixture of the requisite amine Ia (1 equiv) and the appropriate isocyanate (1.2 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was stirred overnight. The mixture was treated with MP-TsOH and rotated for 3 hours. The solid was collected by filtration and washed with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>. The solid was rotated with 2 M NH<sub>3</sub> in MeOH for 2 hours.  
20 The solid was collected by filtration and washed with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was concentrated to give the purified product.

General Procedure L (Parallel Synthesis of Ureas from Anilines using Phoxime Resin)

A mixture of the appropriate aniline (3 equiv) and Phoxime resin (1 equiv) in  
25 CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was rotated for 3 hours. If the aniline had not dissolved, triethylamine (3.5 equiv) was added. The mixture was rotated overnight. The solid was collected by filtration and washed with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, CH<sub>2</sub>Cl<sub>2</sub>, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>. A mixture of the solid and the requisite amine Ia (1.1 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 ml) and toluene (1.5 ml) were heated at 80 °C with shaking overnight and allowed to cool to room temperature. The solid was  
30 collected by filtration and washed with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was concentrated to give the product.

General Procedure M (Parallel Synthesis of Ureas from Anilines using Triphosgene)

A mixture of the appropriate aniline (1.2 equiv), triphosgene (0.4 equiv), and triethylamine (1.4 equiv) in CH<sub>2</sub>Cl<sub>2</sub> was heated at 35 °C for 1 hour. After cooling to room  
35 temperature, the requisite amine Ia (1 equiv) was added. The mixture was stirred

overnight, washed with H<sub>2</sub>O and brine, passed through Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give crude product which was purified by parallel chromatography.

General Procedure N (Parallel Synthesis of Thioureas from Thioisocyanates)

A mixture of the requisite amine Ia (1 equiv) and the appropriate thioisocyanate (1.2 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was stirred overnight. The mixture was treated with MP-TsOH and rotated for 3 hours. The solid was collected by filtration and washed with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>. The solid was rotated with 2 M NH<sub>3</sub> in MeOH for 2 hours. The solid was collected by filtration and washed with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, and CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was concentrated to give the purified product.

General Procedure O (Parallel Synthesis of Carbamates)

A mixture of the requisite amine Ia (1 equiv) and the appropriate succinimide (1.5 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was stirred overnight. If the reaction was not complete, it was heated at 38 °C for 1 hour. The mixture was washed with H<sub>2</sub>O and brine, passed through Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the crude product which was purified via parallel purification (step gradient 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>, 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>).

EXPERIMENTAL SECTION

General

Unless otherwise noted, all non-aqueous reactions were run under a nitrogen atmosphere and Na<sub>2</sub>SO<sub>4</sub> was used to dry all organic layers. Purifications were typically carried out by flash chromatography on silica gel (230-400 mesh) or preparative TLC on Uniplate Silica Gel GF PLC Plates (20 x 20 cm, 1000 microns) from Analtech, Inc., Newark, DE. Alumina used was basic with 6 wt % H<sub>2</sub>O (Brockmann III). Melting points taken in capillary tubes are uncorrected. IR spectra were determined in KBr. NMR spectra were run in CDCl<sub>3</sub>, unless otherwise indicated. <sup>1</sup>H NMR spectra were recorded on 300 MHz instruments and <sup>13</sup>C NMR spectra were recorded at 75.5 MHz. Mass spectral analyses were accomplished using electrospray ionization. Analytical reverse-phase HPLC was performed on Shimadzu system equipped with a diode array spectrometer (range 190-300 nm; Hewlett Packard). The stationary phase was a Zorbax SB-Phenyl Rapid Resolution column (4.6 mm x 50 mm; Hewlett Packard), mobile phase A was 0.1% trifluoroacetic acid, and mobile phase B was CH<sub>3</sub>CN. A flow rate of 2.5 ml/min with a linear gradient of 20-55% B in 5 min and then 55-20% B in 5 min was employed. Other physical and analytical data were obtained by the physical and analytical chemistry group at Roche Bioscience. All parallel synthesis reactions were run in sealed tubes that were vented prior to rotated overnight. Amberlite IRA67 (Aldrich Chemical Co., Milwaukee, Wis., USA) was washed consecutively with CH<sub>2</sub>Cl<sub>2</sub>, MeOH, CH<sub>2</sub>Cl<sub>2</sub>, MeOH, CH<sub>2</sub>Cl<sub>2</sub> and then dried under




vacuum prior to use. All products derived from parallel synthesis reactions were characterized via HPLC-MS.

## EXAMPLES


- 5           The following Preparations (1-6) are useful for preparing synthetic intermediates that can be used to prepare compounds of the invention, as described in the following Examples.

Preparation 1: Preparation of ( $\pm$ )-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine




10

Step A: Preparation of ( $\pm$ )-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexanol



- 15           Following Procedure A, 4-(4-chlorobenzyl)-piperidine (see Preparation 7) (52 mg, 0.25 mmol) was alkylated with 7-oxa-bicyclo[4.1.0]heptane (0.25 ml, 2.5 mmol) in EtOH (0.5 ml) at 80 °C for 3 days. Chromatography of the crude product with 90:9.5:0.5-80:19:1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (68 mg, 88%) as a tan oil which solidified upon standing as a cream solid: mp 100-101.3 °C; IR 3379, 2929 cm<sup>-1</sup>; <sup>1</sup>H NMR δ 1.05-1.76 (m, 12H), 2.02 (dt, *J* = 2.4, 11.6 Hz, 1H), 2.06-2.20 (m, 2H), 2.49 (d, *J* = 7.0 Hz, 2H), 2.51-2.64 (m, 2H), 2.79 (m, 1H), 3.34 (m, 1H), 4.05 (m, 1H), 7.06 (m, 2H), 7.24 (m, 2H); MS *m/z* 308 (M + H)<sup>+</sup>. Anal. (C<sub>18</sub>H<sub>26</sub>ClNO) C, H, N.
- 20


Step B: Preparation of (±)-trans-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine




5

A solution of (±)-trans-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexanol (390 mg, 1.27 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 ml) at 0 °C was treated successively with Et<sub>3</sub>N (350 µl, 2.53 mmol) and MeSO<sub>2</sub>Cl (194 µl, 2.53 mmol), stirred at 0 °C for 2 hours, and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 10% NH<sub>4</sub>OH. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were washed with brine, dried and concentrated. A solution of the residue in THF (3 ml) and 28-30 wt % NH<sub>4</sub>OH (1.2 ml) was stirred at 70 °C for 24 hours, allowed to cool to room temperature, and partitioned between EtOAc and 1 N NaOH. The aqueous phase was extracted with EtOAc and the extracts were washed with brine, dried and concentrated. Chromatography of the residue on alumina with 1:3 EtOAc:MeOH to 100% MeOH and a subsequent chromatography on alumina with 20:1 hexanes:EtOAc to 100% EtOAc followed by 3:1 EtOAc:MeOH to 100% MeOH gave the product (260 mg, 67%) as a tan oil which solidified upon standing: mp 69.1-70.4 °C; <sup>1</sup>H NMR δ 1.03-1.34 (m, 6H), 1.37-1.52 (m, 1H), 1.57-1.77 (m, 5H), 1.92-2.05 (m, 3H), 2.48 (d, *J* = 7.0 Hz, 2H), 2.45-2.64 (m, 3H), 2.73 (m, 1H), 7.06 (m, 2H), 7.23 (m, 2H); MS *m/z* 307 (M + H)<sup>+</sup>.

20 Preparation 2: Preparation of (±)-trans-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine



Step A: Preparation of (±)-trans-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentanol




25

Following General Procedure A, a solution of 4-(4-chlorobenzyl)-piperidine (17.86 g, 85.05 mmol) and 6-oxa-bicyclo[3.1.0]hexane (50 g, 0.6 mol) in EtOH (170 ml) was stirred

at 95 °C for 40 hours, allowed to cool to room temperature, and concentrated. The residue was crystallized in hot CH<sub>2</sub>Cl<sub>2</sub> (80 ml), the crystallization mixture was concentrated to half the volume, and kept at 0 °C overnight and filtered, and the precipitate was rinsed with cold hexanes to give the product (18.2 g, 73%) as a tan solid. The mother liquors were concentrated to half the volume, diluted with CH<sub>2</sub>Cl<sub>2</sub> and kept at -10 °C for 1 hour, and the precipitate was rinsed with cold CH<sub>2</sub>Cl<sub>2</sub> and hexanes to give additional product (1.8 g, 7%) as a tan solid: mp 104.1-105.5 °C; IR 3436, 2928 cm<sup>-1</sup>; <sup>1</sup>H NMR δ 1.19-1.75 (m, 8H), 1.81-1.99 (m, 4H), 2.06 (dt, *J* = 2.5, 11.7 Hz, 1H), 2.47 (m, 1H), 2.50 (d, *J* = 7.0 Hz, 2H), 2.90 (m, 1H), 3.07 (m, 1H), 4.10 (m, 1H), 7.06 (m, 2H), 7.23 (m, 2H); <sup>13</sup>C NMR δ 21.63, 27.35, 32.01, 32.15, 34.31, 37.87, 42.47, 50.47, 52.97, 75.15, 75.22, 128.27, 130.43, 131.55, 139.04; MS *m/z* 294 (M + H)<sup>+</sup>. Anal. (C<sub>17</sub>H<sub>24</sub>ClNO•0.1H<sub>2</sub>O) C, H, N.

Step B: Preparation of (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine




15


Following General Procedure B, a solution of (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentanol (205 mg, 0.697 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.8 ml) at 0 °C was treated successively with Et<sub>3</sub>N (190 µl, 1.4 mmol) and MeSO<sub>2</sub>Cl (110 µl, 1.4 mmol), stirred at 0 °C for 1 hour, and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 10% NH<sub>4</sub>OH. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated to give 220 mg of an oil. A solution of the residue (110 mg) in dioxane (2 ml) and 28-30 wt % NH<sub>4</sub>OH (0.8 ml) was stirred at 70-80 °C overnight, allowed to cool to room temperature, and concentrated. The residue was partitioned between EtOAc and 1 N NaOH, the aqueous phase was extracted with EtOAc, and the extracts were washed with brine, dried and concentrated. Chromatography of the residue on alumina with 10:1 hexanes:EtOAc to 100% EtOAc followed by 95:4.75:0.25-60:38:2 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (87 mg, 85%) as an oil: <sup>1</sup>H NMR δ 1.18-1.71 (m, 9H), 1.76-2.00 (m, 3H), 2.07 (dt, *J* = 2.4, 11.5 Hz, 1H), 2.31 (m, 1H), 2.50 (d, *J* = 6.9 Hz, 2H), 2.86-2.99 (m, 2H), 3.19 (m, 1H), 7.06 (m, 2H), 7.23 (m, 2H); MS *m/z* 293.2 (M + H)<sup>+</sup>.

25

**Preparation 3: Preparation of (±)-cis-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine**




**Step A: Preparation of (±)-trans-4-nitro-benzenesulfonic acid 2-azido-cyclopentyl ester**



A solution of (±)-trans-2-azido-cyclopentanol (1.27 g, 10.0 mmol) (Zhang, Z. da; Scheffold, R. *Helv. Chim. Acta* 1993, 76, 2602) in CH<sub>2</sub>Cl<sub>2</sub> (14 ml) at 0 °C was treated successively with pyridine (0.88 ml, 10.9 mmol) and 4-nitro-benzenesulfonyl chloride (2.22 g, 10.0 mmol) and allowed to warm to room temperature slowly. The reaction was stirred for 4 days, during which additional pyridine (0.9 ml, 11 mmol) and 4-nitro-benzenesulfonic acid (2.2 g, 10 mmol) was added, and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 1 N HCl. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were washed with saturated NaHCO<sub>3</sub>, dried and concentrated. Chromatography of the residue with 10:1-4:1 hexanes:EtOAc gave the product (2.63 g, 84%) as a yellow oil: <sup>1</sup>H NMR δ 1.61-1.90 (m, 4H), 2.00-2.16 (m, 2H), 3.96 (m, 1H), 4.72 (m, 1H), 8.14 (m, 2H), 8.43 (m, 2H).


**Step B: Preparation of (±)-cis-1-(2-azido-cyclopentyl)-4-(4-chlorobenzyl)-piperidine**



A murky solution of (±)-trans-4-nitro-benzenesulfonic acid 2-azido-cyclopentyl ester (630 mg, 2.0 mmol), 4-(4-chlorobenzyl)-piperidine (420 mg, 2.0 mmol), and Et<sub>3</sub>N (280 μl, 2.0 mmol) in CH<sub>3</sub>CN (4 ml) was stirred at room temperature for 10 days and 65 °C for 2 days, allowed to cool to room temperature, and concentrated. The residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 1 N NaOH, the aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated. Chromatography of the residue with 20:1-1:1


hexanes:EtOAc followed by chromatography with 100% CH<sub>2</sub>Cl<sub>2</sub> to 95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (145 mg, 22%) as a tan oil: <sup>1</sup>H NMR δ 1.32-1.90 (m, 13H), 2.33 (m, 1H), 2.49 (d, *J* = 6.4 Hz, 2H), 2.96 (m, 1H), 3.06 (m, 1H), 4.04 (t, *J* = 4.0 Hz, 1H), 7.06 (m, 2H), 7.23 (m, 2H); MS *m/z* 319.2 (M - H)<sup>-</sup>.

5      Step C: Preparation of (±)-cis-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine




A solution of (±)-cis-1-(2-azido-cyclopentyl)-4-(4-chlorobenzyl)-piperidine (210 mg, 0.65 mmol) in THF (2.5 ml) was treated successively with PPh<sub>3</sub> (514 mg, 1.96 mmol) and H<sub>2</sub>O (141 μl, 7.83 mmol), refluxed for 3.5 hours, allowed to cool to room temperature, and concentrated. Chromatography of the residue with 90:9.5:0.5-75:23.75:1.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (183 mg, 95 %) as a colorless oil which solidified upon standing to a cream solid: mp 69.6-71.3 °C; <sup>1</sup>H NMR δ 1.20-1.35 (m, 2H), 1.43-1.93 (m, 11H), 2.17 (m, 1H), 2.49 (d, *J* = 6.9 Hz, 2H), 2.89-3.02 (m, 2H), 3.34 (t, *J* = 4.4 Hz, 1H), 7.06 (m, 2H), 7.23 (m, 2H); <sup>13</sup>C NMR δ 20.72, 27.08, 32.48, 32.61, 38.32, 42.95, 52.14, 53.09, 53.61, 71.49, 128.63, 130.80, 131.88, 139.58; MS *m/z* 293.2 (M + H)<sup>+</sup>.

Preparation 4: Preparation of (±)-cis-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine



20      Step A: Preparation of (±)-trans-4-nitro-benzenesulfonic acid 2-azido-cyclohexyl ester




A solution of (±)-trans-2-azidocyclohexan-1-ol (11.3 g, 80.0 mmol) (Zhang, Z. da; Scheffold, R. *Helv. Chim. Acta* 1993, 76, 2602] in CH<sub>2</sub>Cl<sub>2</sub> (110 ml) at 0 °C was treated successively with pyridine (14.2 ml, 176 mmol) and 4-nitro-benzenesulfonyl chloride (35.6

g, 160 mmol), allowed to warm to room temperature slowly, stirred for 4 days, and partitioned between  $\text{CH}_2\text{Cl}_2$  and 1 N HCl. The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were washed with saturated  $\text{NaHCO}_3$ , dried and concentrated.


Chromatography of the residue with 10:1-1:1 hexanes:EtOAc gave the product (19 g, 72%)  
 5 as a cream solid:  $^1\text{H}$  NMR  $\delta$  1.19-1.39 (m, 3H), 1.53-1.82 (m, 3H), 2.00-2.10 (m, 1H), 2.26 (m, 1H), 3.36 (m, 1H), 4.35 (ddd,  $J = 4.7, 9.2, 10.8$  Hz, 1H), 8.17 (m, 2H), 8.41 (m, 2H).

Step B: Preparation of ( $\pm$ )-cis-1-(2-azido-cyclohexyl)-4-(4-chlorobenzyl)-piperidine



10 A murky solution of ( $\pm$ )-*trans*-4-nitro-benzenesulfonic acid 2-azido-cyclohexyl ester (1.77 g, 5.41 mmol), 4-(4-chlorobenzyl)-piperidine (1.14 g, 5.43 mmol), and  $\text{Et}_3\text{N}$  (0.75 ml, 5.4 mmol) in  $\text{CH}_3\text{CN}$  (11.2 ml) was stirred at room temperature for 17 hours, 65 °C for 31 hours, and 80 °C for 5 days, allowed to cool to room temperature, and concentrated. The residue was partitioned between  $\text{CH}_2\text{Cl}_2$  and 1 N NaOH, the aqueous phase was  
 15 extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were dried and concentrated. Chromatography of the residue with 98:1.9:0.1-95:4.75:0.25  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  to 100 % MeOH and subsequent chromatography with 10:1 hexanes:EtOAc to 100% EtOAc followed by 95:5 EtOAc:MeOH gave, in order of elution, starting ( $\pm$ )-*trans*-4-nitro-benzenesulfonic acid 2-azidocyclohexyl ester (1.2 g, 68%), desired product (155 mg, 9%), and starting  
 20 4-(4-chlorobenzyl)-piperidine (810 mg, 71%). Product:  $^1\text{H}$  NMR  $\delta$  1.19-1.81 (m, 12H), 1.92-2.08 (m, 3H), 2.22 (m, 1H), 2.48 (d,  $J = 7.0$  Hz, 2H), 3.02 (m, 2H), 4.05 (m, 1H), 7.06 (m, 2H), 7.23 (m, 2H); MS  $m/z$  333.2 ( $\text{M} + \text{H}$ ) $^+$ .

Step C: Preparation of ( $\pm$ )-cis-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine




25


A solution of ( $\pm$ )-cis-1-(2-azido-cyclohexyl)-4-(4-chlorobenzyl)-piperidine (155 mg, 0.463 mmol) in THF (1.8 ml) was treated successively with  $\text{PPh}_3$  (364 mg, 1.39 mmol) and

H<sub>2</sub>O (141 µl, 5.56 mmol), refluxed for 3 hours, allowed to cool to room temperature, and concentrated. Chromatography of the residue with 95:4.75:0.25-75:23.75:1.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (121 mg, 85 %) as a cream solid: <sup>1</sup>H NMR δ 1.14-1.93 (m, 15H), 1.96 (dt, *J* = 11.8, 3.5 Hz, 1H), 2.48 (d, *J* = 7.0 Hz, 2H), 3.03-3.13 (m, 2H), 3.30 (m, 1H), 7.06 (m, 2H), 7.23 (m, 2H); MS *m/z* 307.2 (M + H)<sup>+</sup>.

Preparation 5: Preparation of (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutylamine



Step A: Preparation of (±)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutanone



1,2-Bis(trimethylsilyloxy)cyclobutene (5.0 g, 22 mmol) at 0 °C under Ar was treated dropwise during 15 min with a solution of 4-(4-chlorobenzyl)-piperidine (4.56 g, 21.7 mmol) in MeOH (10.9 ml) and allowed to warm to room temperature. The reaction was stirred over a period of 5 hours, during which additional 1,2-bis(trimethylsilyloxy)-cyclobutene (0.99 g, 4.3 mmol) was added, and concentrated. Chromatography of the residue with 95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (4.8 g, 80 %) as a yellow oil: <sup>1</sup>H NMR δ 1.20-1.35 (m, 2H), 1.43-1.64 (m, 3H), 1.93-2.18 (m, 4H), 2.49 (d, *J* = 6.9 Hz, 2H), 2.64-2.91 (m, 3H), 3.14 (m, 1H), 3.90 (m, 1H), 7.05 (m, 2H), 7.23 (m, 2H); MS *m/z* 278.1 (M + H)<sup>+</sup>.

Step B: Preparation of (±)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutanone *O*-methyl-oxime



A solution of (±)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutanone (1.74 g, 6.26 mmol) and MeONH<sub>2</sub>•HCl (2.63 g, 31.3 mmol) in MeOH (20 ml) was stirred at 65 °C

under Ar for 3 hours, allowed to cool to room temperature, and concentrated. The residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>, the aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub>, and the extracts were dried and concentrated. Chromatography of the residue with 95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (1.5 g, 78 %) as a brown oil and predominantly one stereoisomer: <sup>1</sup>H NMR δ 1.05-1.65 (m, 4.5H), 1.92-2.11 (m, 4H), 2.45-2.65 (m, 3H), 2.73-2.96 (m, 2H), 3.22 (m, 1H), 3.73 (m, 1H), 3.82 (m, 3H), 4.57 (m, 0.5H), 7.06 (m, 2H), 7.23 (m, 2H); MS *m/z* 307.1 (M + H)<sup>+</sup>.

Step C: Preparation of (±)-trans-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutylamine




10

A mixture of NaBH<sub>4</sub> (604 mg, 16.0 mmol) in THF (13 ml) under Ar was treated dropwise with trifluoroacetic acid (1.23 ml, 16.0 mmol), stirred for 5 min, treated dropwise with a solution of (±)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutanone O-methyl-oxime (985 mg, 3.21 mmol) in THF (35 ml), and stirred at room temperature for 5 hours. The mixture was treated carefully with 6 N HCl (1.5 ml) until the pH ~ 2, stirred for 10 min, basified with 8 N NaOH until the pH ~ 10, and partitioned between EtOAc and 1 N NaOH. The aqueous phase was extracted with EtOAc and the extracts were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. A solution of the residue in MeOH (30 ml) and 1 N HCl (3 ml) was stirred at 50 °C for 1 hour and at 75 °C for 5 hours, allowed to cool to room temperature, and concentrated. The residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 1 N NaOH, the aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated. Chromatography of the residue on alumina with 10:1 hexanes:EtOAc to 100% EtOAc followed by 98:1.9:0.1-90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave 400 mg of the product (80% pure by <sup>1</sup>H NMR) as a yellow oil which was used without further purification: <sup>1</sup>H NMR δ 1.19-1.90 (m, 9H), 2.11 (m, 1H), 2.28 (m, 1H), 2.44-2.59 (m, 3H), 2.80 (m, 1H), 3.05 (m, 1H), 3.22 (m, 1H), 7.06 (m, 2H), 7.23 (m, 2H); MS *m/z* 279.2 (M + H)<sup>+</sup>.

Preparation 6: Preparation of (±)-cis-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutylamine


30






A solution of ( $\pm$ )-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutanone O-methyl-oxime (438 mg, 1.43 mmol) in THF (13 ml) under Ar was treated dropwise with 1 M  $\text{BH}_3 \cdot \text{THF}$  complex in THF (8.6 ml, 8.6 mmol) and stirred at room temperature for 3 hours and at 75 °C for 20 hours. The reaction was cooled to 0 °C and treated carefully with 6 N HCl (1 ml) until the pH  $\sim$  2. The THF was evaporated and a solution of the residue in EtOH (9 ml) and 6 N HCl (1 ml) was stirred at 75 °C for 1 hour. It was then allowed to cool to room temperature, basified with 8 N NaOH (4 ml) until the pH  $\sim$  10, diluted with  $\text{H}_2\text{O}$  (5 ml) to dissolve the resulting white precipitate, and concentrated. The residue was partitioned between  $\text{CH}_2\text{Cl}_2$  and 1 N NaOH, the aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$ , and the extracts were dried and concentrated. Chromatography of the residue with 90:9.5:0.5-60:38:2  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave, in order of elution, 70 mg of the desired product (80% pure by  $^1\text{H}$  NMR) as a colorless oil which was used without further purification, 48 mg (12%) of pure desired product as a colorless oil, and 125 mg of a mixture of desired product, stereoisomeric ( $\pm$ )-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutylamine, and an unidentified impurity. Product:  $^1\text{H}$  NMR  $\delta$  1.19-1.70 (m, 8H), 1.89-2.05 (m, 3H), 2.50(d,  $J$  = 6.9 Hz, 2H), 2.56 (m, 1H), 2.78 (m, 2H), 3.44 (m, 1H), 7.06 (m, 2H), 7.23 (m, 2H);  $^{13}\text{C}$  NMR  $\delta$  24.39, 25.56, 31.63, 31.76, 38.01, 42.61, 49.17, 49.63, 51.74, 62.51, 128.25, 130.42, 131.50, 139.16; MS  $m/z$  279.2 ( $M + 1$ ) $^+$ .

#### Preparation 7: Preparation of 4-(4-chlorobenzyl)-piperidine



##### Step A: Preparation of 4-(4-Chloro-benzylidene)-piperidine-1-carboxylic acid *tert*-butyl ester




25

The phosphonium salt (10g) was taken up in THF and placed in an ice bath. The KHMDS (42ml) was added slowly, the ice bath was removed, and the reaction was stirred for 45 minutes at room temperature. The reaction solution was then cooled to -78°C and the ketone (4.2g) was added slowly. The reaction was stirred for 30 minutes, the cooling

bath was removed, and the reaction was stirred overnight at room temperature. The reaction solution was poured into a saturated  $\text{NH}_4\text{Cl}$  (100ml) solution, the layers were separated, the aqueous layer was washed twice with EtOAc, the organic layers were combined, dried ( $\text{MgSO}_4$ ), and concentrated to ~ 40ml. The solution was diluted with  
5 hexane and filtered to remove the majority of the  $\text{Ph}_3\text{PO}$ . Chromatography of the crude product with 20:1-10:1 hexane:EtOAc gave the product as a colorless oil (4.7g).


Step B: Preparation of 4-(4-Chloro-benzyl)-piperidine-1-carboxylic acid *tert*-butyl ester



10 The protected piperidine (10g) was dissolved in EtOAc (100ml), the  $\text{PtO}_2$  was added, and the mixture was stirred rapidly under  $\text{H}_2$  for 3 hours. The mixture was filtered through celite and concentrated. The crude product was taken up in hot hexane, filtered and allowed to crystallize. The product was recrystallized with hot hexane to yield the clean product (8.0g). Additional product was isolated from the mother liquor.


15

Step C: Preparation of 4-(4Chlorobenzyl)piperidine



20 Methanol (400ml) was placed in an ice bath and AcCl (60ml) was added. After the addition was completed the solution was stirred at room temperature for one hour. The protected piperidine (62.8g) was added and the solution was stirred at room temperature overnight. The reaction solution was concentrated to ~70ml (when product first started to precipitate out), diluted with ether (500ml), and the product was collected by filtration (44.9g). An additional 3.1g of the product was collected from the mother liquor.

Example 1: Preparation of (±)-*trans*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexyl}-3-(3,4,5-trimethoxyphenyl)urea.




5

Following General Procedure C, (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine (56 mg, 0.18 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (76 mg, 0.36 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) and DMF (1 ml) at room temperature for 1.5 hour. After the combined extracts were washed with H<sub>2</sub>O, dried and concentrated, preparative TLC with 95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH followed by 90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH and subsequent preparative TLC with 93:6.65:0.35 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (52 mg, 55%) as a grey solid: mp 196.9-200.0 °C; IR 1670, 1606, 1545, 1505 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO, 87 °C] δ 1.00-1.25 (m, 6H), 1.38-1.63 (m, 4H), 1.70 (m, 1H), 1.78 (m, 1H), 2.06 (m, 1H), 2.13-2.27 (m, 2H), 2.43 (d, *J* = 6.8 Hz, 2H), 2.44 (m, 1H), 2.59 (m, 1H), 2.74 (m, 1H), 3.39 (m, 1H), 3.63 (s, 3H), 3.73 (s, 6H), 5.69 (d, *J* = 5.3 Hz, 1H), 6.75 (s, 2H), 7.12 (m, 2H), 7.24 (m, 2H), 8.28 (s, 1H); MS *m/z* 516 (M + H)<sup>+</sup>.

15

Example 2: Preparation of (±)-*trans*-N-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexyl}-4-methanesulfonyl-benzamide



20


Following General Procedure E, (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine (92 mg, 0.30 mmol) and 4-methanesulfonyl-benzoic acid (72 mg, 0.36 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) using HOBt (8 mg, 0.06 mmol) and DEC (86 mg, 0.45 mmol) at room temperature for 16 hours. Purification of the crude product by preparative TLC with 93:6.65:0.35 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (105 mg, 72%) as a tan solid: mp 184.5-186.7 °C; IR 1637 cm<sup>-1</sup>; <sup>1</sup>H NMR δ 0.94 (ddd, *J* = 3.9, 11.7, 23.9 Hz, 1H), 1.09-1.93 (m, 12H), 2.05 (dt, *J* = 2.4, 11.5 Hz, 1H), 2.35-2.74 (m, 6H), 3.10 (m, 3H), 3.55-3.66 (m, 1H), 7.04 (m, 2H), 7.16 (m, 1H), 7.23 (m, 2H), 7.95 (m, 2H),

25

8.04 (m, 2H); MS  $m/z$  489 ( $M + H$ )<sup>+</sup>. Anal. (C<sub>26</sub>H<sub>33</sub>ClN<sub>2</sub>O<sub>3</sub>S) H, N; C: calcd, 63.85; found, 60.83. HPLC purity: 99.4%.

**Example 3: Preparation of (±)-*trans*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexyl}-3-(3-methanesulfonyl-phenyl)urea hydrochloride**


5



A solution of triphosgene (60 mg, 0.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) was treated with a solution of 3-methanesulfonyl-phenylamine (120 mg, 0.7 mmol) and Et<sub>3</sub>N (125 μl, 0.90 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml), stirred at 40 °C for 1.5 hours, allowed to cool to room temperature, and added in 2 portions during 30 min to a solution of (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine (88 mg, 0.29 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 ml). The reaction was stirred for 1 hour and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated. Purification of the residue by preparative TLC with 95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (140 mg, 0.28 mmol) as a cream solid. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.5 ml, 0.5 mmol) and EtOAc, allowed to stand at room temperature for 3 days, at 0 °C overnight and at -20 °C overnight, and filtered to give the product (80 mg, 50%) as a white solid: mp 150.2-151.6 °C; MS  $m/z$  504 ( $M + H$ )<sup>+</sup>.

20

**Example 4: Preparation of (±)-*trans*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride**




25

Following General Procedure D, (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine (250 mg, 0.85 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (355 mg, 1.70 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (5 ml) at room temperature for 1 hour. After the combined extracts were dried (MgSO<sub>4</sub>) and concentrated, chromatography of the residue with 95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:Et<sub>3</sub>N and subsequent chromatography with 98:1.9:0.1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:Et<sub>3</sub>N gave the free base (248 mg, 0.494 mmol) as a white foam. A solution

30


of the free base (100 mg, 0.2 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (0.5 ml, 0.5 mmol) and concentrated to give the product (104 mg, 56%) as a tan powder: mp 136.1-138.0 °C; IR 1690, 1607, 1555, 1507  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]  $\delta$  1.40-2.15 (m, 11H), 2.53 (m, 1.6H), 2.70 (m, 0.4H), 2.80-2.95 (m, 2H), 3.33-3.60 (m, 3H), 3.59 (s, 3H), 3.71 (s, 6H), 4.24-4.38 (m, 1H), 6.74 (s, 1.6H), 6.77 (s, 0.4H), 6.85 (d,  $J = 8.2$  Hz, 0.8H), 6.91 (m, 0.2H), 7.21 (m, 2H), 7.33 (m, 2H), 8.77 (s, 0.8H), 8.86 (s, 0.2H), 10.10-10.23 (m, 0.8H), 10.45 (m, 0.2H);  $^{13}\text{C}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]  $\delta$  21.82, 26.88, 29.06, 32.57, 34.82, 40.68, 50.73, 51.01, 51.82, 56.02, 60.52, 71.47, 96.26, 128.51, 131.07, 131.27, 132.61, 136.25, 138.62, 153.10, 155.10; MS  $m/z$  502 ( $\text{M} + \text{H}$ ) $^+$ .

10 **Example 5: Preparation of ( $\pm$ )-*trans*-1-[4-[4-(4-chlorobenzyl)piperidin-1-yl]-tetrahydrofuran-3-yl]-3-(3,4,5-trimethoxyphenyl)urea hydrochloride**



15 Following General Procedure D, ( $\pm$ )-*trans*-4-[4-(4-chlorobenzyl)piperidin-1-yl]-tetrahydrofuran-3-ylamine (3.03 g, 10.3 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (2.37 g, 11.3 mol) were coupled in  $\text{CH}_2\text{Cl}_2$  (60 ml) at 0 °C for 1 hour. Chromatography of the crude product with 1:1 hexanes:EtOAc to 100% EtOAc followed by 50:0.95:0.05-10:0.95:0.05  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the free base (4.62 g, 9.2 mmol) as a white foam. A solution of the free base (3.62 g, 7.2 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (10 ml, 10 mmol) and concentrated to give the product (3.61 g, 83%) as a white solid: mp 229.2-230.9 °C; IR 3273 (br), 2937, 1690, 1606, 1554, 1507  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]  $\delta$  1.54-1.74 (m, 7H), 2.52 (m, 2H), 2.97 (m, 2H), 3.49-3.71 (m, 11H), 4.06 (m, 3H), 4.59 (m, 1H), 6.74 (s, 2H), 7.00 (d,  $J = 7.6$  Hz, 1H), 7.22 (m, 2H), 7.35 (m, 2H), 8.85 (s, 1H), 10.8 (br s, 1H); MS  $m/z$  504 ( $\text{M} + \text{H}$ ) $^+$ .


Step A: Preparation of ( $\pm$ )-*trans*-4-[4-(4-chlorobenzyl)piperidin-1-yl]-tetrahydrofuran-3-ol



Following General Procedure A, 4-(4-chlorobenzyl)-piperidine (10.1 g, 48 mmol) was alkylated with 3,6-dioxabicyclo[3.1.0]hexane (24.7 g, 288 mol) (Barili, P. L.; Berti, G.; Mastorilli, E.; *Tetrahedron* 1993, 49, 6263) in EtOH (75 ml) at 90-95 °C for 45 hours.

- 5 Chromatography of the crude product with CH<sub>2</sub>Cl<sub>2</sub> followed by 99:0.95:0.05-95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (10.7g, 76%) as a white solid: <sup>1</sup>H NMR δ 1.21 (m, 2H), 1.51 (m, 1H), 1.62 (m, 2H), 2.03 (tt, *J* = 2.5, 11.6 Hz, 2H), 2.19 (br, 1H), 2.50 (d, *J* = 6.9 Hz, 2H), 2.73 (m, 2H), 3.08 (m, 1H), 3.61 (dd, *J* = 6.9, 9.3 Hz, 1H), 3.70 (dd, *J* = 3.1, 10.0 Hz, 1H), 3.93 (dd, *J* = 5.7, 10.0 Hz, 1H), 4.05 (dd, *J* = 7.0, 9.3 Hz, 1H), 4.33 (dt, *J* = 3.0, 5.7 Hz, 1H), 7.05 (m, 2H), 7.24 (m, 2H); MS *m/z* 296.2 (M + H)<sup>+</sup>.


Step B: Preparation of (±)-*trans*-4-[4-(4-chlorobenzyl)piperidin-1-yl]-tetrahydrofuran-3-ylamine



15

- Following General Procedure B, (±)-*trans*-4-[4-(4-chlorobenzyl)piperidin-1-yl]-tetrahydrofuran-3-ol (10.65 g, 36 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (150 ml) was treated with Et<sub>3</sub>N (10.2 ml, 72 mmol) and MeSO<sub>2</sub>Cl (5.53 ml, 72 mol) for 1.25 hours and the resultant product was heated in dioxane (205 ml) and NH<sub>4</sub>OH (83 ml) for 4 hours. Chromatography of the
- 20 crude product with 100:1.9:0.1-100:19:1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (9.58 g, 90%) as a yellow oil: <sup>1</sup>H NMR δ 1.21-1.68 (m, 7H), 2.01 (m, 2H), 2.50 (d, *J* = 6.9 Hz, 2H), 2.58 (dt, *J* = 3.4, 6.8 Hz, 2H), 2.72 (m, 1H), 3.06 (m, 1H), 3.50 (m, 2H), 3.64 (dd, *J* = 6.5, 9.3 Hz, 1H), 3.99 (m, 2H), 7.05 (m, 2H), 7.25 (m, 2H); MS *m/z* 295.2 (M + H)<sup>+</sup>.


- Example 6: Preparation of (±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxy-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea.
- 25



A solution of (±)-1-[(1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilyloxy)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea (605

mg, 0.96 mmol) in 1% HCl/EtOH (80 ml) was stirred at room temperature overnight. The EtOH was evaporated and the residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were washed with brine, dried (MgSO<sub>4</sub>) and concentrated. Chromatography of the residue with  
 5 95:4.75:0.25-90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (440 mg, 89%) as a yellow solid: mp 98.7-102.0 °C; MS *m/z* 518 (M + H)<sup>+</sup>.


Step A: Preparation of (±)-(1*R*,2*R*,4*R*)-4-(*tert*-butyl-dimethylsilanyloxy)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentanol



10

Following General Procedure A, 4-(4-chlorobenzyl)-piperidine (741 mg, 3.53 mmol) was alkylated with *cis-tert*-butyl-dimethyl-(6-oxa-bicyclo[3.1.0]hex-3-yloxy)-silane (1.51 g, 7.06 mmol) (Asami, M. *Bull. Chem. Soc. Jpn.* 1990, 63, 1402) in EtOH (5 ml) at  
 15 90-95 °C for 82 hours. Chromatography of the crude product with CH<sub>2</sub>Cl<sub>2</sub> followed by 99:0.95:0.05-95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (1.16 g, 78%) as a yellow foam: <sup>1</sup>H NMR δ 0.00 (s, 6H), 0.81 (s, 9H), 1.19-1.99 (m, 12H), 2.44 (m, 2H), 2.80 (m, 2H), 3.13 (m, 1H), 4.07 (m, 1H), 4.28 (m, 1H), 6.99 (m, 2H), 7.17 (m, 2H); <sup>13</sup>C NMR  
 20 δ -4.55, -4.50, 18.3, 26.1, 32.1, 32.2, 38.1, 39.8, 42.3, 44.0, 52.1, 53.0, 74.0, 75.7, 75.8, 128.7, 130.8, 131.9, 139.2; MS *m/z* 424.2 (M + H)<sup>+</sup>. HPLC purity 98.5%.

Step B: Preparation of (±)-(1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilanyloxy)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine




25

Following General Procedure B, (±)-(1*R*,2*R*,4*R*)-4-(*tert*-butyl-dimethylsilanyloxy)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentanol (1.14 g, 2.68 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 ml) was treated with Et<sub>3</sub>N (740 µl, 5.36 mmol) and MeSO<sub>2</sub>Cl (410 µl, 5.36 mmol) for 2 hours

and the resultant product was heated in dioxane (15.6 ml) and  $\text{NH}_4\text{OH}$  (6.2 ml) for 6 hours. Chromatography of the crude product with 98:1.9:0.1-80:19:1  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the product (620 mg, 55%) as a colorless oil:  $^1\text{H}$  NMR  $\delta$  0.00 (s, 6H), 0.83 (s, 9H), 1.15-1.32 (m, 2H), 1.40-2.11 (m, 11H), 2.45 (m, 2H), 2.60 (m, 1H), 2.80 (m, 1H), 3.02 (m, 1H), 3.14 (m, 1H), 4.20 (m, 1H), 7.02 (m, 2H), 7.19 (m, 2H);  $^{13}\text{C}$  NMR  $\delta$  -4.85, -4.87, 17.9, 25.8, 32.0, 32.1, 37.8, 39.2, 42.4, 44.8, 51.0, 52.8, 54.8, 72.8, 75.8, 128.2, 130.4, 131.5, 139.0; MS  $m/z$  423.2 ( $\text{M} + \text{H}$ ) $^+$ . HPLC purity 99.2%.

Step C: Preparation of ( $\pm$ )-1-[(1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilyloxy)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea




10

Following General Procedure C, ( $\pm$ )-1-[(1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilyloxy)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine (610 mg, 1.44 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (603 mg, 2.88 mmol) were coupled in  $\text{CH}_2\text{Cl}_2$  (8.5 ml) at 0 °C for 1 hour. Chromatography of the crude product with 1:1 hexanes:EtOAc to 100% EtOAc gave the product (665 mg, 73%) as a yellow solid: mp 81.0-84.0 °C; IR 3379 (br), 2929, 1653, 1608, 1555, 1507  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]  $\delta$  0.03 (d, 6H), 0.86 (s, 9H), 1.12-1.69 (m, 8H), 1.95 (m, 2H), 2.18 (m, 1H), 2.46 (m, 2H), 2.79 (m, 3H), 3.58 (s, 3H), 3.71 (s, 6H), 3.93 (m, 1H), 4.16 (m, 1H), 5.94 (d,  $J = 7.9$  Hz, 1H), 6.70 (s, 2H), 7.16 (m, 2H), 7.30 (m, 2H), 8.43 (s, 1H); MS  $m/z$  632 ( $\text{M} + \text{H}$ ) $^+$ .

20

Example 7: Preparation of ( $\pm$ )-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxycyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.




25

A solution of ( $\pm$ )-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxy-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea (102 mg, 0.20 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (0.5 ml, 0.5 mmol) and concentrated to give the product (110 mg, 100%) as a yellow solid: mp 137.0-143.0 °C; IR 3405 (br), 2935, 1685, 1606, 1554,




1507  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $[(\text{CD}_3)_2\text{SO}]$   $\delta$  1.48-2.12 (m, 11H), 2.54 (m, 2H), 2.90 (m, 2H), 3.45 (m, 1H), 3.58 (s, 3H), 3.71 (s, 6H), 4.22 (m, 1H), 4.48 (m, 1H), 5.16 (br s, 1H), 6.42 (d,  $J$  = 9.4 Hz, 1H), 6.70 (s, 2H), 7.21 (m, 2H), 7.34 (m, 2H), 10.19 (br s, 1H); MS  $m/z$  518 ( $M + \text{H}$ ) $^+$ .

- 5 **Example 8: Preparation of ( $\pm$ )-1-((1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxymethyl-cyclopentyl)-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.**




- A solution of ( $\pm$ )-1-((1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilyloxymethyl)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl)-3-(3,4,5-trimethoxyphenyl)urea (1.13 g, 1.75 mmol) in 1% HCl/EtOH (120 ml) was stirred at room temperature for 1 hour. The EtOH was evaporated and the residue was partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were washed with brine, dried ( $\text{MgSO}_4$ ) and concentrated. Chromatography of the residue with
- 15 95:4.75:0.25-90:9.5:0.5  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the free base ( $\pm$ )-1-((1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxymethyl-cyclopentyl)-3-(3,4,5-trimethoxyphenyl)urea (690 mg, 1.30 mmol) as a yellow solid. A solution of the free base (105 mg, 0.20 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (0.5 ml, 0.5 mmol) and concentrated to give the product (112 mg, 74%) as
- 20 a yellow solid: mp 138.0-143.0  $^\circ\text{C}$ ; IR 3417 (br), 2936, 1687, 1606, 1554, 1507  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $[(\text{CD}_3)_2\text{SO}]$   $\delta$  1.28-2.12 (m, 10H), 2.54 (m, 2H), 2.91 (m, 2H), 3.26-3.59 (m, 8H), 3.71 (s, 6H), 4.34 (m, 1H), 4.70 (br s, 1H), 6.66 (d,  $J$  = 8.3 Hz, 1H), 6.73 (s, 2H), 7.21 (m, 2H), 7.32 (m, 2H), 8.78 (s, 1H), 10.13 (br s, 1H); MS  $m/z$  532 ( $M + \text{H}$ ) $^+$ .

- 25 Step A: Preparation of ( $\pm$ )-1-((1*R*,2*R*,4*R*)-4-(*tert*-butyl-dimethylsilyloxymethyl)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentanol.




Following General Procedure A, 4-(4-chlorobenzyl)-piperidine (946 mg, 4.50 mmol) was alkylated with *cis-tert*-butyl-dimethyl-(6-oxa-bicyclo[3.1.0]hex-3-ylmethoxy)silane (1.13 g, 4.96 mmol) (Asami, M. Takahashi, J.; Inoue, S. *Tetrahedron Asymmetry* 1994, 5, 1649) in EtOH (4 ml) at 95 °C for 4.5 days. Chromatography of the crude product with CH<sub>2</sub>Cl<sub>2</sub> followed by 99:0.95:0.05-95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (1.38 g, 74%) as a brown oil: <sup>1</sup>H NMR δ 0.00 (s, 6H), 0.84 (s, 9H), 1.34-1.86 (m, 8H), 2.08-2.31 (m, 5H), 2.48 (m, 2H), 2.80 (m, 1H), 3.13 (m, 1H), 3.28 (m, 1H), 3.48 (m, 2H), 4.28 (m, 1H), 7.00 (m, 2H), 7.19 (m, 2H); MS *m/z* 438.2 (M + H)<sup>+</sup>.

10 Step B: Preparation of (±)-(1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilanyloxymethyl)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine



Following General Procedure B, a solution of (±)-(1*R*,2*R*,4*R*)-4-(*tert*-butyl-dimethylsilanyloxymethyl)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentanol (1.36 g, 3.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15 ml) was treated with Et<sub>3</sub>N (0.83 ml, 6.0 mmol) and MeSO<sub>2</sub>Cl (0.46 ml, 6.0 mmol) for 1 hour and the resultant product was heated in dioxane (17.2 ml) and NH<sub>4</sub>OH (6.9 ml) for 2.5 hours. Chromatography of the crude product with 98:1.9:0.1-90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (780 mg, 57%) as a yellow oil: <sup>1</sup>H NMR δ 0.00 (s, 6H), 0.85 (s, 9H), 0.92-1.70 (m, 11H), 1.96-2.21 (m, 3H), 2.44 (m, 3H), 2.83 (m, 2H), 3.12 (m, 1H), 3.43 (d, *J* = 5.8 Hz, 2H), 7.02 (m, 2H), 7.19 (m, 2H); MS *m/z* 438.2 (M + H)<sup>+</sup>. HPLC purity: 96.4%


25 Step C: Preparation of (±)-1-[(1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilanyloxymethyl)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea



Following General Procedure C, (±)-(1*R*,2*R*,4*S*)-4-(*tert*-butyl-dimethylsilanyloxymethyl)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine (770 mg, 1.76 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (442 mg, 2.11 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (10

ml) at 0 °C for 30 min to give 1.19 g of the product as a yellow solid which was used directly in the next step: MS  $m/z$  646.2 (M + H)<sup>+</sup>.


**Example 9: Preparation of (±)-*trans*-1-[4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl]-3-(3,4,5-trimethoxyphenyl)urea dihydrochloride.**



5

- A solution of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)-ureido]-pyrrolidine-1-carboxylic acid *tert*-butyl ester (410 mg, 0.68 mmol) in 2% HCl/MeOH (70 ml) was stirred at room temperature overnight.
- 10 The MeOH was evaporated and the residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were washed with brine, dried (MgSO<sub>4</sub>) and concentrated. Chromatography of the residue on alumina with CH<sub>2</sub>Cl<sub>2</sub> followed by 50:0.95:0.05-5:0.95:0.05 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (±)-*trans*-1-[4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl]-
- 15 3-(3,4,5-trimethoxyphenyl)urea (225 mg, 0.45 mmol) as a white solid. A solution of the free base (25 mg, 0.05 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.15 ml, 0.15 mmol) and concentrated to give the product (27 mg, 68%) as a yellow solid: mp 157.0-170.0 °C; IR 3416 (br), 2935, 1686, 1606, 1554, 1507 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO] δ 1.50-1.81 (m, 5H), 2.60 (d, *J* = 6.6 Hz, 2H), 3.00-3.99 (m, 20H), 4.68 (m, 1H), 6.76 (s, 2H),
- 20 7.21 (m, 2H), 7.32 (m, 2H); MS  $m/z$  504 (M + H)<sup>+</sup>.

Step A: Preparation of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester




25

Following General Procedure A, 4-(4-chlorobenzyl)-piperidine (1.72 g, 8.19 mmol) was alkylated with 6-oxa-3-aza-bicyclo[3.1.0]hexane-3-carboxylic acid *tert*-butyl ester (1.82 g, 9.82 mmol) [Okada, T.; Sato, H.; Tsuji, T.; Tsushima, T.; Nakai, H.; Yoshida, T.; Matsuura, S. *Chem. Pharm. Bull. Jpn.* 1993, 41, 132] in EtOH (5.5 ml) at 95 °C for 60


hours. Chromatography of the crude product with CH<sub>2</sub>Cl<sub>2</sub> followed by 100:0.95:0.05-30:0.95:0.05 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (1.85 g, 54%) as a yellow solid: mp 61.9-64.5 °C; IR 3421 (br) cm<sup>-1</sup>; <sup>1</sup>H NMR δ 1.26-1.69 (m, 14H), 1.94 (br s, 1H), 2.19 (m, 2H), 2.53 (d, *J* = 6.9 Hz, 2H), 2.89 (m, 2H), 3.05-3.33 (m, 3H), 3.60-3.79 (m, 2H), 4.33 (m, 1H), 7.08 (m, 2H), 7.28 (m, 2H); MS *m/z* 395.2 (M + H)<sup>+</sup>.

Step B: Preparation of (±)-*trans*-3-amino-4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester



Following General Procedure B, a solution of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.80 g, 4.56 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) was treated with Et<sub>3</sub>N (1.26 ml, 9.12 mmol) and MeSO<sub>2</sub>Cl (0.70 ml, 9.12 mmol) for 1 hour and the resultant product was heated in dioxane (26.0 ml) and NH<sub>4</sub>OH (10.5 ml) for 6 hours to give 1.91 g of the product (91% pure by HPLC) as a yellow oil which was used directly for the next step: <sup>1</sup>H NMR δ 1.21-1.65 (m, 17H), 2.17 (m, 2H), 2.49 (d, *J* = 7.0 Hz, 2H), 2.73-3.43 (m, 7H), 7.05 (m, 2H), 7.25 (m, 2H); MS *m/z* 394.2 (M + H)<sup>+</sup>.

Step C: Preparation of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)ureido]-pyrrolidine-1-carboxylic acid *tert*-butyl ester




20

Following General Procedure C, (±)-*trans*-3-amino-4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester (394 mg, ~ 0.9 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (250 mg, 1.2 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (6.0 ml) at 0 °C for 1 hour. Chromatography of the crude product with 1:1 hexanes:EtOAc to 100% EtOAc gave the product (445 mg, 77% from (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester) as a white solid: mp 99.0-102.5 °C; IR 3371 (br), 2932, 1655, 1607, 1552, 1507 cm<sup>-1</sup>; <sup>1</sup>H NMR δ 1.17-1.80

25


(m, 14H), 2.18 (m, 2H), 2.48 (d,  $J = 7.0$  Hz, 2H), 2.95-3.51 (m, 8H), 3.81 (m, 9H), 4.32 (br, 1H), 5.30 (br, 1H), 6.61 (s, 2H), 7.02 (m, 2H), 7.20 (m, 2H); MS  $m/z$  603.2 ( $M + H$ )<sup>+</sup>.

Example 10: Preparation of (±)-*cis*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.




Following General Procedure D, a solution of (±)-*cis*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine (110 mg, 0.38 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) at 0 °C was treated with 5-isocyanato-1,2,3-trimethoxybenzene (98 mg, 0.47 mmol), stirred for 1 hour, and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated. Chromatography of the residue with EtOAc followed by 90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH and a subsequent chromatography with 1:3 hexanes:EtOAc to 100% EtOAc followed by 95:4.75:0.25 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (190 mg, 0.38 mmol) as a colorless oil. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was treated with 1 N HCl in Et<sub>2</sub>O (1 ml, 1 mmol) and concentrated to give the product (193 mg, 96%) as a white solid: mp 117.5-122.5 °C; IR 1692, 1606, 1557, 1507 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO] δ 1.33-2.15 (m, 11H), 2.51 (m, 2H), 2.64 (br d,  $J = 7.8$  Hz, 0.5H), 2.80-2.95 (m, 1.5H), 3.15-3.90 (m, 12H), 4.41 (m, 1H), 6.74 (s, 2H), 7.08-7.24 (m, 3H), 7.34 (m, 2H), 9.01 (br s, 0.7H), 9.06 (br s, 0.3H), 9.25-9.47 (m, 1H); HRMS (FAB) calcd for C<sub>27</sub>H<sub>37</sub>ClN<sub>3</sub>O<sub>4</sub> 502.2473 ( $M + H$ )<sup>+</sup>, found 502.2471.

Example 11: Preparation of (±)-*cis*-N-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-4-methanesulfonyl-benzamide hydrochloride.




Following General Procedure F, ( $\pm$ )-*cis*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine (105 mg, 0.358 mmol) and 4-methanesulfonyl-benzoic acid (86 mg, 0.43 mmol) were coupled in  $\text{CH}_2\text{Cl}_2$  (2 ml) using HOBt (10 mg, 0.07 mmol) and DEC (138 mg, 0.719 mmol) at 0 °C for 2.5 hours. Chromatography of the crude product with  $\text{CH}_2\text{Cl}_2$  followed by 98:1.9:0.1  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the free base (176 mg, 0.36 mmol) as a colorless oil. A solution of the free base in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (0.8 ml, 0.8 mmol) and concentrated to give the product (183 mg, 100%) as a cream foam: mp 125.3-131.9 °C; IR 1660  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$  1.63-2.23 (m, 11H), 2.56 (d,  $J$  = 6.4 Hz, 2H), 2.56-2.85 (m, 2H), 3.05 (s, 0.3H), 3.06 (s, 2.7H), 3.20-3.34 (m, 1H), 3.60 (br d,  $J$  = 11.7 Hz, 1H), 3.78 (br d,  $J$  = 10.8 Hz, 1H), 4.93 (m, 1H), 7.00-7.11 (m, 2H), 7.25 (m, 2H), 8.03 (m, 2H), 8.50 (m, 2H), 8.95 (m, 0.1H), 9.06 (br d,  $J$  = 8.3 Hz, 1H), 11.65-11.90 (m, 1H);  $^{13}\text{C}$  NMR  $\delta$  21.06, 26.85, 29.18, 29.38, 33.31, 36.77, 41.57, 44.76, 50.69, 53.67, 54.39, 68.88, 127.87, 129.08, 129.85, 130.64, 132.72, 137.61, 138.20, 143.57, 166.50; HRMS (FAB) calcd for  $\text{C}_{25}\text{H}_{31}\text{ClN}_2\text{O}_3\text{S}$  475.1822 ( $\text{M} + \text{H}$ ) $^+$ , found 475.1823.

Example 12: Preparation of ( $\pm$ )-*trans*-1-{1-acetyl-4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.




A solution of ( $\pm$ )-*trans*-1-[4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl]-3-(3,4,5-trimethoxyphenyl)urea (80 mg, 0.16 mmol) and  $\text{Et}_3\text{N}$  (26  $\mu\text{l}$ , 0.19 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml) at 0 °C was treated dropwise with acetic anhydride (15  $\mu\text{l}$ , 0.16 mmol), stirred at 0 °C for 10 min, and partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were dried and concentrated. Purification of the residue by preparative TLC with 10:0.95:0.05  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the free base as a white solid. A solution of the free base in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (0.2 ml, 0.2 mmol) and concentrated to give the product (84 mg, 91%) as a tan solid: mp 155.0-161.0 °C; IR 3415 (br), 2936, 1691, 1608, 1554, 1508  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]  $\delta$  1.56-1.97 (m, 8H), 2.50-3.94 (m, 20H), 4.70 (m, 1H), 6.76 (s, 2H), 6.98 (m, 1H), 7.21 (m, 2H), 7.32 (m, 2H), 8.81 (s, 1H), 10.72 (s, 1H); MS  $m/z$  545 ( $\text{M} + \text{H}$ ) $^+$ .

**Example 13: Preparation of (±)-cis-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexyl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.**



- Following General Procedure D, (±)-cis-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine (53 mg, 0.17 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (40 mg, 0.19 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) at 0 °C for 1 hour. Purification of the crude product by preparative TLC with 93:6.65:0.35 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (75 mg, 0.15 mmol) as a tan foam. A solution of the free base (100 mg, 0.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.3 ml, 0.3 mmol) and concentrated to give the product (83 mg, 85%) as a tan foam: mp 120.1-137.3 °C; IR 1691, 1607, 1558, 1507 cm<sup>-1</sup>; <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 1.35-2.15 (m, 13H), 2.58 (d, *J* = 6.4 Hz, 1.8H), 2.72 (m, 0.2H), 2.87-2.98 (m, 2H), 3.22 (m, 1H), 3.59 (m, 1H), 3.70 (s, 0.3H), 3.73 (s, 2.7H), 3.78 (s, 0.6H), 3.82 (s, 5.4H), 3.92 (m, 1H), 4.55 (m, 1H), 6.76 (s, 1.8H), 6.77 (s, 0.2H), 7.17 (m, 2H), 7.27 (m, 2H); MS *m/z* 516 (M + H)<sup>+</sup>. Anal. (C<sub>28</sub>H<sub>39</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>4</sub>•1.15H<sub>2</sub>O) C, H, N.


**Example 14: Preparation of (±)-cis-N-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexyl}-4-methanesulfonyl-benzamide hydrochloride.**



- Following General Procedure F, (±)-cis-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclohexylamine (30 mg, 0.10 mmol) and 4-methanesulfonyl-benzoic acid (24 mg, 0.12 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) using HOBT (3 mg, 0.02 mmol) and DEC (30 mg, 0.16 mmol) at 0 °C for 2 hours and at room temperature for 2 hours. Purification of the crude product by preparative TLC with 90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (46 mg, 0.09 mmol) as a white solid. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.2 ml, 0.2 mmol) and concentrated to give the product (47 mg, 88%) as a cream solid: mp 123.7-149.4 °C; IR 1659 cm<sup>-1</sup>; <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 1.28-2.20 (m, 13H), 2.59 (d, *J* = 6.2 Hz, 2H), 2.90-3.03 (m, 1H), 3.17 (s, 3H), 3.35 (m, 1H), 3.61 (m, 1H), 3.89 (m, 1H), 4.47-4.91 (m, 2H), 7.17 (d, *J* = 8.3 Hz, 2H), 7.28 (m, 2H), 8.10 (m, 4H);

MS  $m/z$  489 ( $M + H$ )<sup>+</sup>. Anal. ( $C_{26}H_{34}Cl_2N_2O_3S$ ); calcd: C, 59.42; H, 6.52; N, 5.33; found: C, 55.39; H, 5.94; N, 4.88. HPLC purity: 98.6%.

**Example 15: Preparation of (±)-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-1-methanesulfonyl-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.**




5

A solution of (±)-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea (78 mg, 0.15 mmol) and pyridine (13  $\mu$ l, 0.15 mmol) in  $CH_2Cl_2$  (2.5 ml) at 0 °C was treated dropwise with  $MeSO_2Cl$  (12  $\mu$ l, 0.15 mmol) and allowed to warm to room temperature slowly. The reaction was stirred for 1 hour, during which additional pyridine (3  $\mu$ l, 0.04 mmol) and  $MeSO_2Cl$  (3  $\mu$ l, 0.04 mmol) was added, and partitioned between  $CH_2Cl_2$  and 10% aqueous NaOH. The aqueous phase was extracted with  $CH_2Cl_2$  and the extracts were washed with brine, dried and concentrated. Purification of the residue by preparative TLC with 10:0.95:0.05  $CH_2Cl_2$ :MeOH: $NH_4OH$  gave the free base as a yellow semi-solid. A solution of the free base in  $CH_2Cl_2$  was treated with 1 N HCl in  $Et_2O$  (0.3 ml, 0.3 mmol) and concentrated to give the product (87 mg, 91%) as a tan solid: mp 148.0-158.0 °C; IR 3405 (br), 2933, 1691, 1608, 1554, 1507  $cm^{-1}$ ;  $^1H$  NMR [ $(CD_3)_2SO$ ]  $\delta$  1.51-1.78 (m, 5H), 2.50 (m, 2H), 3.05-3.71 (m, 21H), 4.66 (m, 1H), 6.74 (s, 2H), 6.91 (br d, 1H), 7.21 (m, 2H), 7.34 (m, 2H), 8.84 (s, 1H), 10.75 (s, 1H); MS  $m/z$  581 ( $M + H$ )<sup>+</sup>. Anal. ( $C_{27}H_{38}Cl_2N_4O_6S \cdot 1.35H_2O$ ) C, H, N.

20

**Example 16: Preparation of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)-ureido]-pyrrolidine-1-sulfonic acid dimethylamide hydrochloride.**



25


A solution of (±)-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea (80 mg, 0.16 mmol) and  $Et_3N$  (33  $\mu$ l, 0.24 mmol) in  $CH_2Cl_2$  (2 ml) was treated with dimethylsulfamoyl chloride (21  $\mu$ l, 0.20 mmol), stirred at room temperature for 20 hours, and partitioned between  $CH_2Cl_2$  and saturated

30




NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated. Purification of the residue by preparative TLC with 10:0.95:0.05 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base as a white solid. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.35 ml, 0.35 mmol) and concentrated to give the product (93 mg, 93%) as a white solid: mp 141.0-143.0 °C; IR 3378 (br), 2937, 1691, 1607, 1555, 1508 cm<sup>-1</sup>; <sup>1</sup>H NMR ([ (CD<sub>3</sub>)<sub>2</sub>SO], D<sub>2</sub>O added, 87 °C) δ 1.53 (m, 2H), 1.86 (m, 3H), 2.58 (d, *J* = 6.7 Hz, 2H), 2.84 (s, 6H), 3.07-3.86 (m, 18H), 4.60 (m, 1H), 6.74 (s, 2H), 7.22 (m, 2H), 7.34 (m, 2H); MS *m/z* 610 (M + H)<sup>+</sup>. Anal. (C<sub>28</sub>H<sub>41</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>6</sub>•1H<sub>2</sub>O) C, H, N.

- 10 Example 17: Preparation of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)-ureido]-pyrrolidine-1-carboxylic acid dimethylamide hydrochloride.



- 15 A solution of (±)-*trans*-1-[4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl]-3-(3,4,5-trimethoxyphenyl)urea (90 mg, 0.18 mmol) and Et<sub>3</sub>N (37 µl, 0.27 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) at 0 °C was treated with dimethylcarbonyl chloride (20 µl, 0.22 mmol), allowed to warm to room temperature slowly, stirred for 75 min, and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated. Purification of the residue by preparative TLC with 6:0.95:0.05 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free amine as a white solid. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.2 ml, 0.2 mmol) and concentrated to give the product (70 mg, 69%) as a yellow solid: mp 142.0-144.5 °C; IR 3272 (br), 2936, 1685, 1608, 1555, 1507 cm<sup>-1</sup>; <sup>1</sup>H NMR ([ (CD<sub>3</sub>)<sub>2</sub>SO], D<sub>2</sub>O added, 87 °C) δ 1.54 (m, 2H), 1.84 (m, 3H), 2.54 (m, 2H), 2.82 (s, 6H), 3.05-3.76 (m, 18H), 4.55 (m, 1H), 6.76 (s, 2H), 7.22 (m, 2H), 7.34 (m, 2H); MS *m/z* 574 (M + H)<sup>+</sup>. Anal. (C<sub>29</sub>H<sub>41</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>5</sub>•1.1H<sub>2</sub>O) C, H, N.

Example 18: Preparation of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)-ureido]-pyrrolidine-1-carboxylic acid amide hydrochloride.




5

A solution of (±)-*trans*-1-[4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl]-3-(3,4,5-trimethoxyphenyl)urea (120 mg, 0.24 mmol) and NaOCN (36 mg, 0.55 mmol) in CH<sub>3</sub>CN (2 ml) was treated with trifluoroacetic acid (36 µl, 0.48 mmol). The reaction mixture was stirred at room temperature for 20 hours and concentrated. The residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>, the aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub>, and the extracts were dried and concentrated. Purification of the residue by preparative TLC with 5:0.95:0.05 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base as a white solid. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.2 ml, 0.2 mmol) and concentrated to give the product (70 mg, 54%) as a yellow solid: mp 158.0-162.0 °C; IR 3390 (br), 2935, 1654, 1605, 1554, 1508 cm<sup>-1</sup>; <sup>1</sup>H NMR ([ (CD<sub>3</sub>)<sub>2</sub>SO], D<sub>2</sub>O added, 87 °C) δ 1.53-1.86 (m, 5H), 2.59 (d, *J* = 6.8 Hz, 2H), 3.07-3.86 (m, 18H), 4.60 (m, 1H), 6.74 (s, 2H), 7.22 (m, 2H), 7.34 (m, 2H); MS *m/z* 546 (M + H)<sup>+</sup>. Anal. (C<sub>27</sub>H<sub>37</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>5</sub>•1.1H<sub>2</sub>O) C, H, N

10

15

**Example 19: Preparation of (±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-methoxymethyl-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.**




5 A mixture of (±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxymethyl-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea (160 mg, 0.30 mmol) and silica gel (1.5 g) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) at 0 °C was treated slowly with a fresh 0.5 N solution of diazomethane in Et<sub>2</sub>O (60 ml, 30 mmol), allowed to warm to room temperature, stirred for 4 hours, and filtered. The silica gel was washed with methanol and the filtrate was

10 concentrated. Purification of the residue by preparative TLC with 100:19:1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH followed by another preparative TLC with 100:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (50 mg, 0.09 mmol) as a white foam. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.3 ml, 0.3 mmol) and concentrated to give the product (48 mg, 30%) as a white solid: mp 109.0-112.0 °C; IR

15 3420 (br), 2934, 1686, 1606, 1554, 1507 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO] δ 1.32-2.28 (m, 10H), 2.54 (m, 2H), 2.91 (m, 2H), 3.26-3.50 (m, 8H), 3.59 (s, 3H), 3.71 (s, 6H), 4.33 (m, 1H), 6.62 (d, *J* = 8.3 Hz, 1H), 6.77 (s, 2H), 7.20 (m, 2H), 7.34 (m, 2H), 8.75 (s, 1H), 10.14 (br s, 1H); MS *m/z* 546 (M + H)<sup>+</sup>. Anal. (C<sub>29</sub>H<sub>41</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>5</sub>•0.9H<sub>2</sub>O) C, H, N.

**Example 20: Preparation of (±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-methoxy-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.**

20




A mixture of (±)-1-[(1*R*,2*R*,4*S*)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-4-hydroxy-cyclopentyl]-3-(3,4,5-trimethoxyphenyl)urea (105 mg, 0.20 mmol) and silica

25 gel (1.0 g) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) at 0 °C was treated slowly with a fresh 0.5 N solution of diazomethane in Et<sub>2</sub>O (45 ml, 22.5 mmol), allowed to warm to room temperature slowly, stirred for 4 hours, and filtered. The silica gel was washed with methanol and the filtrate was concentrated. Purification of the residue by preparative TLC with 100:19:1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (43 mg, 0.08 mmol) as a yellow solid. A

30 solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.3 ml, 0.3 mmol)


and concentrated to give the product (46 mg, 40%) as a tan solid: mp 112.5-119.5 °C; IR 3390 (br), 2935, 1686, 1607, 1555, 1506  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $[(\text{CD}_3)_2\text{SO}]$   $\delta$  1.46-1.98 (m, 7H), 2.22 (m, 2H), 2.54 (m, 2H), 2.90 (m, 2H), 3.24 (s, 3H), 3.42 (m, 2H), 3.58-3.85 (m, 11H), 4.50 (m, 1H), 6.36 (d,  $J = 9.1$  Hz, 1H), 6.70 (s, 2H), 7.21 (m, 2H), 7.35 (m, 2H), 10.10 (s, 1H); MS  $m/z$  532 ( $\text{M} + \text{H}$ ) $^+$ . Anal. ( $\text{C}_{28}\text{H}_{39}\text{Cl}_2\text{N}_3\text{O}_5 \cdot 0.7\text{H}_2\text{O}$ ) C, H, N.

**Example 21: Preparation of ( $\pm$ )-*trans*-2-{3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)-ureido]-pyrrolidin-1-yl}-acetamide dihydrochloride.**



A solution of ( $\pm$ )-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea (90 mg, 0.18 mmol) in anhydrous DMF (2.5 ml) was treated successively with *i*-Pr<sub>2</sub>NEt (47  $\mu$ l, 0.27 mmol) and 2-iodoacetamide (40 mg, 0.22 mmol), stirred at room temperature for 24 hours, and partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$ , and the extracts were washed with  $\text{H}_2\text{O}$  and brine, dried and concentrated. Purification of the residue by preparative TLC with 10:0.95:0.05  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the free amine as a yellow solid. A solution of the free base in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (0.3 ml, 0.3 mmol) and concentrated to give the product (54 mg, 50%) as a tan solid: mp 160.0-165.0 °C; IR 3414 (br), 2933, 1691, 1607, 1557, 1507  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $[(\text{CD}_3)_2\text{SO}]$   $\delta$  1.55-1.77 (m, 5H), 2.49 (m, 2H), 3.00-3.71 (m, 20H), 4.72 (br s, 1H), 6.76 (s, 2H), 7.01 (br d, 1H), 7.21 (m, 2H), 7.34 (m, 2H), 7.58 (br s, 1H), 7.83 (br s, 1H), 9.12 (br s, 1H); MS  $m/z$  560 ( $\text{M} + \text{H}$ ) $^+$ . Anal. ( $\text{C}_{28}\text{H}_{40}\text{Cl}_3\text{N}_5\text{O}_5 \cdot 1.3\text{H}_2\text{O}$ ) C, H, N.

**Example 22: Preparation of ( $\pm$ )-*trans*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(4-methoxy-naphthalen-2-yl)urea.**




25

A solution of triphosgene (47 mg, 0.16 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 ml) was treated dropwise with a solution of ( $\pm$ )-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine (140 mg, 0.48 mmol) and *N,N*-diisopropylethylamine (180  $\mu$ l, 1.0 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml) and added to a solution of 4-methoxy-naphthalen-2-ylamine (83 mg, 0.48 mmol) and

30


*N,N*-diisopropylethylamine (180  $\mu$ l, 1.0 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml). The reaction was stirred for 4 hours, washed with 1 N  $\text{KHSO}_4$ , saturated  $\text{NaHCO}_3$ , and brine, dried, and concentrated. Purification of the residue by preparative TLC with 100:9.5:0.5  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the product (100 mg, 43%) as a yellow solid: mp 175-190  $^\circ\text{C}$ ;  
 5 MS  $m/z$  492 ( $\text{M} + \text{H}$ ) $^+$ .

**Example 23: Preparation of ( $\pm$ )-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea dihydrochloride.**



A solution of ( $\pm$ )-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea (100 mg, 0.20 mmol) in MeOH (1 ml) was treated with methyl vinyl sulfone (18  $\mu$ l, 0.21 mmol). The mixture was stirred at room temperature overnight and concentrated. Purification of the residue by preparative TLC with 7:0.95:0.05  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the free amine as a white solid. A solution of  
 15 the free base in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in  $\text{Et}_2\text{O}$  (0.5 ml, 0.5 mmol) and concentrated to give the product (110 mg, 83%) as a white solid: mp 210.2-211.8  $^\circ\text{C}$ ; IR 3422 (br), 2930, 1686, 1607, 1556, 1507  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]  $\delta$  1.54-1.77 (m, 5H), 2.51 (m, 2H), 3.05-3.71 (m, 25H), 4.71 (br s, 1H), 6.75 (s, 2H), 6.92 (br d, 1H), 7.21 (m, 2H), 7.34 (m, 2H), 9.01 (s, 1H); MS  $m/z$  609 ( $\text{M} + \text{H}$ ) $^+$ . Anal. ( $\text{C}_{29}\text{H}_{43}\text{Cl}_3\text{N}_4\text{O}_6\text{S} \cdot 1.15\text{H}_2\text{O}$ )  
 20 C, H, N.


**Example 24: Preparation of ( $\pm$ )-*trans*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-[2-(3,4,5-trimethoxyphenyl)-ethyl]urea hydrochloride.**



A solution of triphosgene (47 mg, 0.16 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 ml) was treated dropwise with a solution of ( $\pm$ )-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentylamine (140 mg, 0.48 mmol) and *N,N*-diisopropylethylamine (180  $\mu$ l, 1.0 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml) and added to a solution of 2-(3,4,5-trimethoxyphenyl)-ethan-1-amine (101 mg, 0.48 mmol) and *N,N*-diisopropylethylamine (180  $\mu$ l, 1.0 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml). The reaction was stirred

overnight and washed with saturated  $\text{NaHCO}_3$ . The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were washed with brine, dried, and concentrated. Purification of the residue by preparative TLC with 10:1  $\text{CH}_2\text{Cl}_2$ :MeOH gave the free base (83 mg, 0.16 mmol) as a yellow oil. A solution of the free base in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in Et<sub>2</sub>O (0.3 ml, 0.3 mmol) and concentrated to give the product (86 mg, 32%) as a yellow solid: mp 90.3-95 °C; MS  $m/z$  530 ( $M + H$ )<sup>+</sup>.


**Example 25: Preparation of (±)-*trans*-3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)-ureido]-pyrrolidine-1-sulfonic acid amide hydrochloride.**



A solution of chlorosulfonyl isocyanate (23  $\mu\text{l}$ , 0.26 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml) was treated with *t*-BuOH (21  $\mu\text{l}$ , 0.22 mmol), stirred at room temperature for 1 hour, and treated with a solution of (±)-*trans*-1-[4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl]-3-(3,4,5-trimethoxyphenyl)urea (110 mg, 0.22 mmol) and pyridine (19  $\mu\text{l}$ , 0.24 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 ml). The reaction was stirred at room temperature for 2 days during which additional chlorosulfonyl isocyanate (46  $\mu\text{l}$ , 0.52 mmol) and *t*-BuOH (42  $\mu\text{l}$ , 0.44 mmol) was added, and partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were dried and concentrated. Purification of the residue by preparative TLC with 10:0.95:0.05  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave 55 mg of a yellow solid.


A solution of the solid in 10% HCl/MeOH (20 ml) was stirred at room temperature overnight. The MeOH was evaporated and the residue was partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated  $\text{NaHCO}_3$ . The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were dried ( $\text{MgSO}_4$ ) and concentrated. Purification of the residue by preparative TLC with 5:0.95:0.05  $\text{CH}_2\text{Cl}_2$ :MeOH: $\text{NH}_4\text{OH}$  gave the free base as a yellow solid. A solution of the free base in  $\text{CH}_2\text{Cl}_2$  was treated with 1 N HCl in Et<sub>2</sub>O (0.1 ml, 0.1 mmol) and concentrated to give the product (20 mg, 48%) as a tan solid: mp 155.7-160.0 °C; IR 3404 (br), 1686, 1607, 1554, 1507  $\text{cm}^{-1}$ ; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]  $\delta$  1.48-1.77 (m, 5H), 2.54 (m, 2H), 3.00-3.83 (m, 18H), 4.60 (m, 1H), 6.73 (s, 2H), 6.85 (m, 1H), 7.11 (br s, 2H), 7.21 (m, 2H), 7.35 (m, 2H), 8.76 (s, 1H), 10.41 (s, 1H); MS  $m/z$  582 ( $M + H$ )<sup>+</sup>. Anal. (C<sub>26</sub>H<sub>37</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>6</sub>S•1.1H<sub>2</sub>O) C, H, N.

**Example 26: Preparation of (±)-*trans*-N-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutyl}-4-methanesulfonyl-benzamide hydrochloride.**



Following General Procedure F, a solution of (±)-*trans*-2-[4-(4-chlorobenzyl)-  
 5 piperidin-1-yl]-cyclobutylamine (130 mg, 80% pure, ~0.38 mmol) and  
 4-methanesulfonyl-benzoic acid (115 mg, 0.575 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) at 0 °C was  
 treated successively with 1-hydroxybenzotriazole hydrate (HOBt) (13 mg, 0.10 mmol) and  
 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (DEC) (138 mg, 0.719  
 mmol), allowed to warm to room temperature slowly, stirred for 3 days, and partitioned  
 10 between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub>  
 and the extracts were dried and concentrated. Purification of the residue by preparative  
 TLC with 90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (118 mg, 0.26 mmol) as a  
 white solid. A solution of the free base (93 mg, 0.20 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N  
 HCl in Et<sub>2</sub>O (0.5 ml, 0.5 mmol) and concentrated to give the product (90 mg, 21% from  
 15 (±)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutanone *O*-methyl-oxime) as a tan  
 powder: mp 173.2-184.1 °C; IR 1657 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO] δ 1.45-2.20 (m, 9H), 2.52  
 (m, 2H), 2.65-2.80 (m, 2H), 3.20-3.45 (m, 5H), 3.68-3.79 (m, 1H), 4.79-4.92 (m, 1H), 7.21  
 (m, 2H), 7.34 (m, 2H), 8.00-8.19 (m, 4H), 9.38 (d, *J* = 8.3 Hz, 1H), 11.04-11.30 (m, 1H);  
<sup>13</sup>C NMR [(CD<sub>3</sub>)<sub>2</sub>SO] δ 17.86, 21.23, 28.00, 34.70, 40.50, 43.21, 46.54, 48.69, 49.15, 64.74,  
 20 126.90, 128.07, 128.34, 130.60, 130.78, 138.20, 138.35, 143.08, 164.24; MS *m/z* 461 (*M* +  
 H)<sup>+</sup>. Anal. (C<sub>24</sub>H<sub>30</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>3</sub>S•0.4 CH<sub>2</sub>Cl<sub>2</sub>) C, H, N.


**Example 27: Preparation of (±)-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-1,1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-yl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride**



Following General Procedure C, (±)-*trans*-4-[4-(4-chlorobenzyl)piperidin-1-yl]-  
 25 1,1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-ylamine (60 mg, 0.18 mmol) and  
 5-isocyanato-1,2,3-trimethoxybenzene (44 mg, 0.21 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (1.5


ml) at 0 °C for 1 hour. Purification of the crude product by preparative TLC with 90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH and a subsequent preparative TLC with EtOAc gave the free base (55 mg, 0.10 mmol) as yellow solid. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.3 ml, 0.3 mmol) and concentrated to give the product (59 mg, 57%) as a yellow solid: mp 162.0-166.0 °C; IR 3377 (br), 2935, 1690, 1606, 1556, 1506 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO] δ 1.48 (m, 2H), 1.78 (m, 3H), 2.56 (d, *J* = 6.5 Hz, 2H), 2.89 (m, 2H), 3.21-3.60 (m, 6H), 3.64 (s, 3H), 3.74 (s, 6H), 3.99 (m, 1H), 4.78 (m, 1H), 6.74 (s, 2H), 7.19 (m, 2H), 7.31 (m, 2H); MS *m/z* 552 (M + H)<sup>+</sup>. Anal. (C<sub>27</sub>H<sub>37</sub>Cl<sub>2</sub>N<sub>3</sub>O•0.4H<sub>2</sub>O) C, H, N.

10      Step A: Preparation of (±)-trans-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-1.1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-yl}-carbamic acid ethyl ester



A solution of (4-chloro-1.1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-yl)-carbamic acid ethyl ester (500 mg, 2.07 mmol) (Ohba, K.; Mori, K.; Kitahara, T.; Kitamura, S.; Matsui, M. *Agr. Biol. Chem.* 1974, 38, 1679) and 4-(4-chlorobenzyl)-piperidine (599 mg, 2.85 mmol) in EtOH (5 ml) was refluxed for 22 hours, allowed to cool to room temperature, and concentrated. Chromatography of the residue on silica gel with 95:4.75:0.25-90:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH and a subsequent preparative TLC with 10:1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH gave the product (171 mg, 20%) as a yellow solid: <sup>1</sup>H NMR δ 1.16-1.33 (m, 5H), 1.45-1.68 (m, 3H), 2.03 (m, 1H), 2.26 (m, 1H), 2.50 (d, *J* = 7.0 Hz, 2H), 2.80 (m, 2H), 2.92-3.08 (m, 2H), 3.19-3.36 (m, 2H), 3.78 (m, 1H), 4.14 (q, *J* = 7.1 Hz, 2H), 4.25 (m, 1H), 5.24 (br, 1H), 7.04 (m, 2H), 7.21 (m, 2H); MS *m/z* 415.2 (M + H)<sup>+</sup>.

25      Step B: Preparation of (±)-trans-4-[4-(4-chlorobenzyl)piperidin-1-yl]-1.1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-ylamine




A solution of (±)-trans-4-[4-(4-chlorobenzyl)piperidin-1-yl]-1.1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-yl}-carbamic acid ethyl ester (170 mg, 0.41 mmol) in 48%



aqueous HBr (3 ml) was refluxed for 16 hours, allowed to cool to room temperature, and poured onto a mixture of ice and granular Na<sub>2</sub>CO<sub>3</sub>. The reaction mixture was completely neutralized and extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were washed with brine, dried and concentrated. Purification of the residue by preparative TLC with 100:9.5:0.5

- 5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (60 mg, 43%) as a yellow solid: <sup>1</sup>H NMR δ 1.26 (m, 2H), 1.49 (m, 1H), 1.66 (m, 4H), 2.00 (m, 1H), 2.29 (m, 1H), 2.50 (d, *J* = 7.0 Hz, 2H), 2.77-2.91 (m, 3H), 3.00-3.22 (m, 3H), 3.44 (m, 1H), 3.65 (m, 1H), 7.05 (m, 2H), 7.26 (m, 2H); MS *m/z* 343.1 (M + H)<sup>+</sup>.


- Example 28: Preparation of (±)-*trans*-3-{3-[4-(4-chlorobenzyl)piperidin-1-yl]-4-[3-(3,4,5-trimethoxyphenyl)-ureido]-pyrrolidin-1-yl}-3-oxo-propionic acid.



- A solution of (±)-*trans*-1-{4-[4-(4-chlorobenzyl)piperidin-1-yl]-pyrrolidin-3-yl}-3-(3,4,5-trimethoxyphenyl)urea (82 mg, 0.16 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) at 0 °C was treated successively with Et<sub>3</sub>N (27 µl, 0.20 mmol) and methyl malonyl chloride (19 µl, 0.18 mmol) and allowed to warm to room temperature slowly. The reaction was stirred for 19 hours, during which additional Et<sub>3</sub>N (27 µl, 0.20 mmol) and methyl malonyl chloride (19 µl, 0.18 mmol) was added, and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated to give 87 mg of ester as a yellow solid which was used directly in the next step: <sup>1</sup>H NMR δ 1.08-1.61 (m, 5H), 2.03-2.48 (m, 4H), 2.74-3.93 (m, 21H), 4.50 (m, 1H), 5.82 (m, 1H), 6.70 (s, 2H), 7.02 (m, 2H), 7.25 (m, 2H), 7.49 (br s, 1H); MS *m/z* 603.2 (M + H)<sup>+</sup>.


- A solution of the ester in 5% KOH/MeOH (6 ml) was stirred at room temperature overnight. The MeOH was evaporated, the residue was treated carefully with 1 N HCl until the pH ~ 7 and extracted with CH<sub>2</sub>Cl<sub>2</sub>, and extracts were dried and concentrated. Purification of the residue by preparative TLC with 95:5 MeOH:NH<sub>4</sub>OH and a subsequent preparative TLC with 4:0.95:0.05 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the product (40 mg, 42%) as a yellow solid: mp 184.0-184.9 °C; IR 3377, 2924, 1607, 1554, 1506 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO] δ 1.15-1.49 (m, 5H), 2.15 (m, 2H), 2.50 (m, 2H), 3.00-3.71 (m, 18H), 4.62 (m, 1H), 6.80 (s, 2H), 7.13 (m, 3H), 7.25 (m, 2H), 8.87 (m, 1H); MS *m/z* 589 (M + H)<sup>+</sup>.

Example 29: Preparation of (±)-*trans*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutyl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.



A solution of (±)-*trans*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutylamine (130 mg, 80% pure, ~0.38 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) at 0 °C was treated with  
 5 5-isocyanato-1,2,3-trimethoxybenzene (110 mg, 0.53 mmol). The reaction was stirred at 0 °C for 2 hours, during which additional 5-isocyanato-1,2,3-trimethoxybenzene (25 mg, 0.12 mmol) was added, and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extracts were dried and concentrated. Chromatography of the residue with 1:1 hexanes:EtOAc to 100% EtOAc followed by  
 10 95:4.75:0.25-50:47.5:2.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (138 mg, 0.28 mmol) as a glassy solid. A solution of the free base (74 mg, 0.15 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.5 ml, 0.5 mmol) and concentrated to give the product (76 mg, 25% from (±)-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutanone *O*-methyl-oxime) as a yellow powder: mp 127.4-128.2 °C; IR 1691, 1607, 1555, 1507 cm<sup>-1</sup>; <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO, 77 °C] δ  
 15 1.45-1.85 (m, 6H), 1.93-2.20 (m, 3H), 2.54 (d, *J* = 6.7 Hz, 2H), 2.65-2.80 (m, 2H), 3.07 (br s, 0.5H), 3.28 (m, 1H), 3.47-3.62 (m, 1.5H), 3.62 (s, 3H), 3.72 (s, 6H), 4.45-4.58 (m, 1H), 6.75 (m, 2H), 6.90-7.03 (m, 1H), 7.19 (m, 2H), 7.31 (m, 2H), 8.67 (br s, 0.8H), 8.76 (br s, 0.2H), 10.70-11.00 (m, 1H); MS *m/z* 488 (M + H)<sup>+</sup>. Anal. (C<sub>26</sub>H<sub>34</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>4</sub>•0.65H<sub>2</sub>O) C, H, N.


20 **Example 30: Preparation of (±)-*cis*-N-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutyl}-4-methanesulfonyl-benzamide hydrochloride.**



Following General Procedure F, (±)-*cis*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutylamine (100 mg, 90% pure, ~0.33 mmol) and 4-methanesulfonyl-benzoic acid  
 25 (88 mg, 0.44 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) using HOBt (25 mg, 0.18 mmol) and DEC (106 mg, 0.55 mmol) at 0 °C for 3.5 hours. Purification of the crude product by preparative TLC with 10:0.95:0.05 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH gave the free base (102 mg, 0.22 mmol) as a colorless oil. A solution of the free base (82 mg, 0.18 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.5 ml, 0.5 mmol) and concentrated to give the product (83

mg, 64%) as a white solid: mp 130.0-132.5 °C; IR 3438, 2924, 1662, 1541 cm<sup>-1</sup>; MS *m/z* 461 (M + H)<sup>+</sup>. Anal. (C<sub>24</sub>H<sub>30</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>3</sub>S•0.35H<sub>2</sub>O) C, H, N.

**Example 31: Preparation of (±)-*cis*-1-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutyl}-3-(3,4,5-trimethoxyphenyl)urea hydrochloride.**



5

Following General Procedure D, (±)-*cis*-2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclobutylamine (40 mg, 0.15 mmol) and 5-isocyanato-1,2,3-trimethoxybenzene (37 mg, 0.18 mmol) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) at 0 °C for 1 hour. Purification of the crude product by preparative TLC with 100:9.5:0.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH and a subsequent preparative TLC with EtOAc gave the free base as a yellow solid. A solution of the free base in CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 N HCl in Et<sub>2</sub>O (0.1 ml, 0.1 mmol) and concentrated to give the product (26 mg, 34%) as a tan solid: mp 123.7-127.9 °C; IR 3423, 2925, 1690, 1607, 1556, 1507. Anal. (C<sub>26</sub>H<sub>35</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>4</sub>•0.7H<sub>2</sub>O) C, H, N.

15

**Example 32: The following compound was prepared using General Procedure C, with the appropriate amine and isocyanate.**

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3,4,5-trimethoxyphenyl)urea.

20

**Example 33: The following compounds were prepared using General Procedure E, with the appropriate amine and carboxylic acid.**

(±)-*trans*-N-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]cyclohexyl}-2-[5-(3,4-dimethoxy-phenyl)-pyrimidin-2-ylsulfanyl]acetamide; and

25

(±)-*trans*-N-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]cyclopentyl}-2-[5-(3,4-dimethoxy-phenyl)-pyrimidin-2-ylsulfanyl]acetamide.

**Example 34: The following compounds were prepared following General Procedure F, using the appropriate amine and carboxylic acid.**

30

(±)-*cis*-N-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-2-[5-(3,4-dimethoxy-phenyl)-pyrimidin-2-ylsulfanyl]-acetamide hydrochloride;

(±)-*cis*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclohexyl}-2-[5-(3,4-dimethoxy-phenyl)-pyrimidin-2-ylsulfanyl]-acetamide hydrochloride; and

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclobutyl}-2-[5-(3,4-dimethoxy-phenyl)-pyrimidin-2-ylsulfanyl]-acetamide hydrochloride.

**Example 35:** The following compounds were prepared following General Procedure G, using the appropriate amine and sulfonyl chloride.

(±)-*trans*-4-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-fluoro-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-4-ethyl-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-methoxy-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-2,4-difluoro-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-trifluoromethyl-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-4-methyl-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-4-fluoro-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-2-fluoro-benzenesulfonamide;

(±)-*trans*-2-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzenesulfonamide;

(±)-*trans*-4-Acetyl-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzenesulfonamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-4-nitro-benzenesulfonamide;

- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-nitro-benzenesulfonamide;
- (±)-*trans*-3-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-2-methyl-benzenesulfonamide;
- 5 (±)-*trans*-Naphthalene-1-sulfonic acid {2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-amide;
- (±)-*trans*-2-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-4-fluoro-benzenesulfonamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-10 3,4-dimethoxy-benzenesulfonamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-2,5-dimethoxy-benzenesulfonamide;
- (±)-*trans*-2,3-Dichloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzenesulfonamide;
- 15 (±)-*trans*-2,4-Dichloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzenesulfonamide;
- (±)-*trans*-3,4-Dichloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzenesulfonamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-20 4-methanesulfonyl-benzenesulfonamide;
- (±)-*trans*-4-Bromo-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzenesulfonamide;
- (±)-*trans*-4-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-nitro-benzenesulfonamide;
- 25 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-2-nitro-benzenesulfonamide; and
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-2-nitro-4-trifluoromethyl-benzenesulfonamide.
- 30 **Example 36:** The following compounds were prepared following General Procedure H, using the appropriate amine and acid chloride.
- (±)-*trans*-4-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-benzamide;

- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-fluoro-benzamide;
- (±)-*trans*-2-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide;
- 5 (±)-*trans*-3-Chloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-methyl-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
10 3-methyl-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-methyl-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-methoxy-benzamide;
- 15 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-trifluoromethyl-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-methoxy-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
20 4-fluoro-benzamide;
- (±)-*trans*-3-Bromo-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide;
- (±)-*trans*-2,4-Dichloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide;
- 25 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-trifluoromethyl-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-trifluoromethyl-benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
30 3,5-bis-trifluoromethyl-benzamide;
- (±)-*trans*-4-*tert*-Butyl-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-ethoxy-benzamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-cyano-benzamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-cyano-benzamide;

5 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-methoxy-benzamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-nitro-benzamide;

10 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-nitro-benzamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3,5-dimethoxy-benzamide;

(±)-*trans*-3,4-Dichloro-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide;

15 (±)-*trans*-4-Bromo-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide; and

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-iodo-benzamide.

20 **Example 37:** The following compounds were prepared following General Procedure I,  
using the appropriate amine and carboxylic acid.

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-isopropyl-benzamide;

25 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-isophthalamide  
acid methyl ester;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
4-methanesulfonyl-benzamide;

(±)-*trans*-3-Acetyl-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
benzamide;

30 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-*o*-tolyl-acetamide;

(±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-*m*-tolyl-acetamide;

- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-*p*-tolyl-acetamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-(4-fluoro-phenyl)-acetamide;
- 5 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-(3-methoxy-phenyl)-acetamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-(4-methoxy-phenyl)-acetamide;
- 10 (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-(3-chloro-phenyl)-acetamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
2-(4-chloro-phenyl)-acetamide;
- (±)-*trans*-5-Methyl-1H-indole-2-carboxylic acid {2-[4-(4-chlorobenzyl)piperidin-  
1-yl]-cyclopentyl}-amide;
- 15 (±)-*trans*-5-Fluoro-1H-indole-2-carboxylic acid {2-[4-(4-chlorobenzyl)piperidin-  
1-yl]-cyclopentyl}-amide;
- (±)-*trans*-2-(3-Bromo-phenyl)-*N*-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-  
cyclopentyl}-acetamide;
- (±)-*trans*-*N*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
20 2-(3,4,5-trimethoxyphenyl)-acetamide;
- (±)-*trans*-1H-Indole-2-carboxylic acid {2-[4-(4-chlorobenzyl)piperidin-1-yl]-  
cyclopentyl}-amide;
- (±)-*trans*-5-Methoxy-1H-indole-2-carboxylic acid {2-[4-(4-chlorobenzyl)-  
piperidin-1-yl]-cyclopentyl}-amide; and
- 25 (±)-*trans*-5-Chloro-1H-indole-2-carboxylic acid {2-[4-(4-chlorobenzyl)piperidin-  
1-yl]-cyclopentyl}-amide.

**Example 38:** The following compounds were prepared following General Procedure J, using the appropriate amine and isocyanate.

- 30 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-chloro-phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-fluoro-phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-*o*-tolyl-urea;



- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(2-chloro-phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3-chloro-phenyl)urea;
- 5 (±)-*trans*-1-(3-Bromo-phenyl)-3-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-  
cyclopentyl}-urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-*p*-tolyl-urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-*m*-tolyl-urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(4-methoxy--  
10 phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3-methoxy-phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(2-methoxy-phenyl)urea;
- 15 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-cyclohexyl-urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3-trifluoromethyl-phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-trifluoromethyl-phenyl)urea;
- 20 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-ethoxy-phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-isopropyl-phenyl)urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
25 3-(4-cyano-phenyl)urea;
- (±)-*trans*-1-(3-Acetyl-phenyl)-3-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-  
cyclopentyl}-urea;
- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3-nitro-phenyl)urea;
- 30 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-nitro-phenyl)urea;
- (±)-*trans*-2-(3-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
ureido)-benzoic acid methyl ester;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3,5-dimethoxy-phenyl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(2,4-dichloro-phenyl)urea;

5 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3,5-dichloro-phenyl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3,4-dichloro-phenyl)urea;

10 (±)-*trans*-1-(4-Bromo-phenyl)-3-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-  
cyclopentyl}-urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(2-fluoro-phenyl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(3-fluoro-phenyl)urea;

15 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(2,4-difluoro-phenyl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(2,3-dichloro-phenyl)urea;

20 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-ethyl-phenyl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-naphthalen-1-yl-urea;

(±)-*trans*-1-(3,5-Bis-trifluoromethyl-phenyl)-3-{2-[4-(4-chlorobenzyl)piperidin-  
1-yl]-cyclopentyl}-urea;

25 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(2-nitro-phenyl)urea; and

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
3-(4-iodo-phenyl)urea.

30 **Example 39:** The following compounds were prepared following General Procedure K,  
using the appropriate amine and isocyanate.

(±)-*trans*-1-(4-Acetyl-phenyl)-3-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-  
cyclopentyl}-urea;

- (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-phenyl-urea;  
 (±)-*trans*-1-Benzyl-3-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(3-chloro-2-methyl-phenyl)urea;  
 5 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-phenethyl-urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-phenethyl-urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(2,5-dimethyl-phenyl)urea;  
 (±)-*trans*-1-[(1*R*,2*R*)-2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl]-  
 10 3-[(*S*)-1-phenyl-ethyl]urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(3,4-dimethoxy-phenyl)urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(2-trifluoromethoxy-phenyl)urea;  
 15 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(4-trifluoromethoxy-phenyl)urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(2-trifluoromethyl-phenyl)urea;  
 (±)-*trans*-1-(4-*tert*-Butyl-phenyl)-3-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-  
 20 cyclopentyl}-urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(2-thiophen-2-yl-ethyl)urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(4-methyl-benzyl)urea;  
 25 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(4-chloro-3-nitro-phenyl)urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(4-fluoro-benzyl)urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 30 3-(2-methyl-benzyl)urea;  
 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-  
 3-(3-methyl-benzyl)urea;

(±)-*trans*-1-(2-Chlorobenzyl)-3-{2-[4-(4-chlorobenzyl)piperidin-1-yl]-cyclopentyl}-urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(4-methoxy-benzyl)urea; and

5 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3,4-dichlorobenzyl)urea.

**Example 40:** The following compounds were prepared following General Procedure L, using the appropriate aniline and Phoxime resin.

10 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3-ethoxy-phenyl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-naphthalen-2-yl-urea;

15 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-isoquinolin-3-yl-urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(2-methylquinolin-6-yl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3-methylamino-phenyl)urea;

20 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-quinolin-3-yl-urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-quinolin-2-yl-urea;

25 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(5-hydroxy-naphthalen-2-yl)urea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(8-hydroxy-quinolin-2-yl)urea; and

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(5,7-dimethyl-[1,8]naphthyridin-2-yl)urea.

30

**Example 41:** The following compounds were prepared following General Procedure M, using the appropriate aniline.

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-quinolin-6-yl-urea;

(±)-*trans*-*N*-[3-(3-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-ureido)-phenyl]-acetamide;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(2,3,4-trimethoxyphenyl)urea; and

5 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3-methanesulfonyl-phenyl)urea.

Example 42: The following compounds were prepared following General Procedure N, using the appropriate thioisocyanate.

10 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-*o*-tolyl-thiourea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-*m*-tolyl-thiourea;

1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(3-chloro-phenyl)-thiourea;

15 (±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-*p*-tolyl-thiourea;

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(4-fluoro-phenyl)-thiourea; and

(±)-*trans*-1-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-3-(4-methoxy-phenyl)-thiourea.

20

Example 43: The following compounds were prepared following General Procedure O, using the appropriate succinimide.

(±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid 4-fluoro-benzyl ester;

25 (±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid 3-chlorobenzyl ester;

(±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid 4-chlorobenzyl ester;

30 (±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid 2-chlorobenzyl ester;

(±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid 3-nitro-benzyl ester;

(±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid  
3-trifluoromethyl-benzyl ester;

(±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid  
3,4-dichlorobenzyl ester;

5 (±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid  
3,5-dichlorobenzyl ester;

(±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid benzyl  
ester; and

(±)-*trans*-{2-[4-(4-Chlorobenzyl)piperidin-1-yl]-cyclopentyl}-carbamic acid 4-  
10 nitro-benzyl ester.

#### Example 44: Formulation Examples

The following are representative pharmaceutical Formulations containing a  
compound of Formula (I).

#### 15 Tablet Formulation

The following ingredients are mixed intimately and pressed into single scored tablets.

Ingredient	Quantity per tablet, mg
compound of this invention	400
20 cornstarch	50
croscarmellose sodium	25
lactose	120
magnesium stearate	5

#### Capsule Formulation

25 The following ingredients are mixed intimately and loaded into a hard-shell gelatin  
capsule.

	Ingredient	Quantity per capsule, mg
	compound of this invention	200
	lactose, spray-dried	148
5	magnesium stearate	2

### Suspension Formulation

The following ingredients are mixed to form a suspension for oral administration.

	Ingredient	Amount
	compound of this invention	1.0 g
10	fumaric acid	0.5 g
	sodium chloride	2.0 g
	methyl paraben	0.15 g
	propyl paraben	0.05 g
	granulated sugar	25.5 g
15	sorbit (70% solution)	12.85 g
	Veegum K (Vanderbilt Co.)	1.0 g
	flavoring	0.035 ml
	colorings	0.5 mg
	distilled water	q.s. to 100 ml

### 20 Injectable Formulation

The following ingredients are mixed to form an injectable Formulation.

Ingredient	Amount
compound of this invention	0.2 g

sodium acetate buffer solution,	0.4M 2.0 ml
HCl (1N) or NaOH (1N)	q.s. to suitable pH
water (distilled, sterile)	q.s. to 20 ml

### Liposomal Formulation

- 5 The following ingredients are mixed to form a liposomal Formulation.

Ingredient	Amount
compound of this invention	10 mg
L-.alpha.-phosphatidylcholine	150 mg
tert-butanol	4 ml

- 10 Freeze dry the sample and lyophilize overnight. Reconstitute the sample with 1 ml 0.9% saline solution. Liposome size can be reduced by sonication.

### Example 45: CCR-3 Receptor Binding Assay--In Vitro

- The CCR-3 antagonistic activity of the compounds of the invention was determined by their ability to inhibit the binding of <sup>125</sup>I eotaxin to CCR-3 L1.2 transfectant cells (see  
 15 Ponath, P. D. et al., J. Exp. Med., Vol. 183, 2437-2448, (1996)).

- The assay was performed in Costar 96-well polypropylene round bottom plates. Test compounds were dissolved in DMSO and then diluted with binding buffer (50 mM HEPES, 1 mM CaCl.sub.2, 5 mM MgCl<sub>2</sub>, 0.5% bovine serum albumin (BSA), 0.02% sodium azide, pH 7.24) such that the final DMSO concentration was 2%. 25 µl of the test  
 20 solution or only buffer with DMSO (control samples) was added to each well, followed by the addition of 25 µl of <sup>125</sup>I-eotaxin (100 pmol) (NEX314, New England Nuclear, Boston, Mass.) and 1.5 x 10<sup>5</sup> of the CCR-3 L1.2 transfected cells in 25 µl binding buffer. The final reaction volume was 75 µl.

- After incubating the reaction mixture for 1 hour at room temperature, the reaction  
 25 was terminated by filtering the reaction mixture through polyethylenimine treated Packard Unifilter GF/C filter plate (Packard, Chicago, Ill.). The filters were washed four times with ice cold wash buffer containing 10 mM HEPES and 0.5M sodium chloride (pH 7.2) and dried at 65°C. for approximately 10 minutes. 25 µl/well of Microscint-20® scintillation



fluid (Packard) was added and the radioactivity retained on the filters was determined by using the Packard TopCount®.

Compounds of this invention were active in this assay.

Compound Number from Table 1	IC50 (μM)
1	0.5574
2	0.0185
3	1.1438
4	0.8644
5	2.6906
6	1.8558
7	1.4841
8	6.0949
9	5.1191
10	0.5122

5    **Example 46: Inhibition of Eotaxin Mediated Chemotaxis of CCR-3 L1.2 Transfectant Cells--In Vitro Assay**

10    The CCR-3 antagonistic activity of the compounds of this invention can be determined by measuring the inhibition of eotaxin mediated chemotaxis of the CCR-3 L1.2 transfectant cells, using a slight modification of the method described in Ponath, P. D. et al., J. Clin. Invest. 97: 604-612 (1996). The assay is performed in a 24-well chemotaxis plate (Costar Corp., Cambridge, Mass.). CCR-3 L1.2 transfectant cells are grown in culture medium containing RPMI 1640, 10% Hyclone® fetal calf serum, 55 mM 2-mercaptoethanol and Geneticin 418 (0.8 mg/ml). 18-24 hours before the assay, the transfected cells are treated with n-butyric acid at a final concentration of 5 mM/1x10<sup>6</sup>

cells/ml, isolated and resuspended at  $1 \times 10^7$  cells/ml in assay medium containing equal parts of RPMI 1640 and Medium 199 (M 199) with 0.5% bovine serum albumin.

Human eotaxin suspended in phosphate buffered saline at 1 mg/ml is added to bottom chamber in a final concentration of 100 nm. Transwell culture inserts (Costar Corp., Cambridge, Mass.) having 3 micron pore size are inserted into each well and L1.2 cells ( $1 \times 10^6$ ) are added to the top chamber in a final volume of 100  $\mu$ l. Test compounds in DMSO are added both to the top and bottom chambers such that the final DMSO volume is 0.5%. The assay is performed against two sets of controls. The positive control contained cells with no test compound in the top chamber and only eotaxin in the lower chamber. The negative control contains cells with no test compound in the top chamber and neither eotaxin nor test compound in lower chamber. The plate is incubated at 37 °C. After 4 hours, the inserts are removed from the chambers and the cells that have migrated to the bottom chamber are counted by pipetting out 500  $\mu$ l of the cell suspension from the lower chamber to 1.2 ml Cluster tubes (Costar) and counting them on a FACS for 30 seconds.

#### 15 Example 47: Inhibition of Eotaxin Mediated Chemotaxis of Human Eosinophils--In Vitro Assay

The ability of compounds of the invention to inhibit eotaxin mediated chemotaxis of human eosinophils can be assessed using a slight modification of procedure described in Carr, M. W. et al., Proc. Natl. Acad. Sci. USA, 91: 3652-3656 (1994). Experiments are performed using 24 well chemotaxis plates (Costar Corp., Cambridge, Mass.). Eosinophils are isolated from blood using the procedure described in PCT Application, Publication No. WO 96/22371. The endothelial cells used are the endothelial cell line ECV 304 obtained from European Collection of Animal Cell Cultures (Porton Down, Salisbury, U.K.). Endothelial cells are cultured on 6.5 mm diameter Biocoat.RTM. Transwell tissue culture inserts (Costar Corp., Cambridge, Mass.) with a 3.0  $\mu$ M pore size. Culture media for ECV 304 cells consists of M199, 10% Fetal Calf Serum, L-glutamine and antibiotics. Assay media consists of equal parts RPMI 1640 and M199, with 0.5% BSA. 24 hours before the assay  $2 \times 10^5$  ECV 304 cells are plated on each insert of the 24-well chemotaxis plate and incubated at 37 °C. 20 nM of eotaxin diluted in assay medium is added to the bottom chamber. The final volume in bottom chamber is 600  $\mu$ l. The endothelial coated tissue culture inserts are inserted into each well.  $10^6$  eosinophil cells suspended in 100  $\mu$ l assay buffer are added to the top chamber. Test compounds dissolved in DMSO are added to both top and bottom chambers such that the final DMSO volume in each well was 0.5%. The assay is performed against two sets of controls. The positive control contains cells in the

top chamber and eotaxin in the lower chamber. The negative control contains cells in the top chamber and only assay buffer in the lower chamber. The plates are incubated at 37 °C. in 5% CO<sub>2</sub> /95% air for 1-1.5 hours.

The cells that migrate to the bottom chamber are counted using flow cytometry. 500  
5 µl of the cell suspension from the lower chamber are placed in a tube, and relative cell counts are obtained by acquiring events for a set time period of 30 seconds.

**Example 48: Inhibition of Eosinophil Influx Into the Lungs of Ovalbumin Sensitized Balb/c Mice by CCR-3 Antagonist--In Vivo Assay**


The ability of the compounds of the invention to inhibit leukocyte infiltration into  
10 the lungs can be determined by measuring the inhibition of eosinophil accumulation into the bronchioalveolar lavage (BAL) fluid of Ovalbumin (OA)-sensitized balb/c mice after antigen challenge by aerosol. Briefly, male balb/c mice weighing 20-25g are sensitized with OA (10 µg in 0.2 ml aluminum hydroxide solution) intraperitoneally on days 1 and 14. After a week, the mice are divided into ten groups. Test compound or only vehicle  
15 (control group) or anti-eotaxin antibody (positive control group) is administered either intraperitoneally, subcutaneously or orally. After 1 hour, the mice are placed in a Plexiglass box and exposed to OA aerosol generated by a PARISTAR.TM. nebulizer (PARI, Richmond, Va.) for 20 minutes. Mice which have not been sensitized or challenged are included as a negative control. After 24 or 72 hours, the mice are anesthetized (urethane,  
20 approx. 1 g/kg, i.p.), a tracheal cannula (PE 60 tubing) is inserted and the lungs are lavaged four times with 0.3 ml PBS. The BAL fluid is transferred into plastic tubes and kept on ice. Total leukocytes in a 20 µl aliquot of the BAL fluid is determined by Coulter Counter.TM. (Coulter, Miami, Fla.). Differential leukocyte counts are made on Cytospin.TM. preparations which have been stained with a modified Wright's stain (DiffQuick.TM.) by  
25 light microscopy using standard morphological criteria.

The foregoing invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding. It will be obvious to one of skill in the art that changes and modifications may be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be  
30 illustrative and not restrictive. The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled.

All patents, patent applications and publications cited in this application are hereby incorporated by reference in their entirety for all purposes to the same extent as if each individual patent, patent application or publication were so individually denoted.

Claims

1. A compound of Formula (I):



(I)

5 wherein:

R<sup>1</sup> is (C<sub>1</sub>-C<sub>2</sub>)alkylene ;

R<sup>2</sup> is optionally substituted phenyl;

R<sup>3</sup> is hydrogen, C<sub>1-6</sub> alkyl, acyl, aryl, or aryl C<sub>1-6</sub> alkyl;

ring A is a C<sub>3-7</sub> cycloalkyl, heterocyclyl, or optionally substituted phenyl;

10 L is -C(=O)-, -C(=S)-, -SO<sub>2</sub>-, -C(=O)N(R<sub>a</sub>)-, -C(=S) N(R<sub>a</sub>)-, -SO<sub>2</sub> N(R<sub>a</sub>)-, -C(=O)O-, -C(=S)O-, -S(=O)<sub>2</sub>O-;

where R<sub>a</sub> is hydrogen, C<sub>1-6</sub> alkyl, acyl, aryl, aryl C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy carbonyl, or benzyloxycarbonyl;

X is absent, -(CR'R'')O-, -(CR'R'')S-, -(CR'R'')NR<sub>b</sub>- or C<sub>1-6</sub> alkylene;

15 where R' and R'' are independently hydrogen or C<sub>1-6</sub> alkyl, and R<sub>b</sub> is hydrogen or C<sub>1-6</sub> alkyl;

R<sup>4</sup> is aryl or heteroaryl; and

R<sup>5</sup> is hydrogen or C<sub>1-6</sub> alkyl;

provided that when R<sup>1</sup> is -CH<sub>2</sub>-, R<sup>2</sup> is phenyl, R<sup>3</sup> is hydrogen, R<sup>5</sup> is hydrogen, A is phenyl, L is -C(=O)NH- and X is absent, then R<sup>4</sup> is not 2,5-difluorophenyl; or

prodrugs, individual isomers, racemic and non-racemic mixtures of isomers, and pharmaceutically acceptable salts thereof;

wherein

the term "optionally substituted phenyl" refers to a phenyl group which is optionally substituted with one or more substituents selected from the group consisting of C<sub>1-6</sub> alkyl, halo C<sub>1-6</sub> alkyl, hydroxy C<sub>1-6</sub> alkyl, heteroalkyl, acyl, acylamino, amino, C<sub>1-6</sub> alkylamino, di  
 5 C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl, -SO<sub>2</sub>NR'R" (where R' and R" are independently hydrogen or C<sub>1-6</sub> alkyl), C<sub>1-6</sub> alkoxy, halo C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkoxycarbonyl, carbamoyl, hydroxy, halo, nitro, cyano, mercapto, methylenedioxy or ethylenedioxy;

the term "acyl" refers to a radical -C(O)R, where R is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl,  
 10 C<sub>3-7</sub> cycloalkyl C<sub>1-6</sub> alkyl, phenyl or phenyl C<sub>1-6</sub> alkyl;

the term "aryl" refers to a monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 10 ring atoms which is optionally substituted with one or more substituents selected from the group consisting of C<sub>1-6</sub> alkyl, halo C<sub>1-6</sub> alkyl, hydroxy C<sub>1-6</sub> alkyl, heteroalkyl, acyl, acylamino, amino, C<sub>1-6</sub> alkylamino, di C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub>  
 15 alkylsulfonyl, -SO<sub>2</sub>NR'R" (where R' and R" are independently hydrogen or C<sub>1-6</sub> alkyl), C<sub>1-6</sub> alkoxy, halo C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkoxycarbonyl, carbamoyl, hydroxy, halo, nitro, cyano, mercapto, methylenedioxy or ethylenedioxy;

the term "heteroaryl" refers to a monocyclic or bicyclic radical of 5 to 12 ring atoms having at least one aromatic ring containing one, two, or three ring heteroatoms selected from N,  
 20 O, or S, the remaining ring atoms being C, with the understanding that the attachment point of the heteroaryl radical will be on an aromatic ring, and the heteroaryl ring is optionally substituted independently with one or more substituents selected from C<sub>1-6</sub> alkyl, halo C<sub>1-6</sub> alkyl, hydroxy C<sub>1-6</sub> alkyl, heteroalkyl, acyl, acylamino, amino, C<sub>1-6</sub> alkylamino, di C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl, -  
 25 SO<sub>2</sub>NR'R" (where R' and R" are independently hydrogen or C<sub>1-6</sub> alkyl), C<sub>1-6</sub> alkoxy, halo C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkoxycarbonyl, carbamoyl, hydroxy, halo, nitro, cyano, mercapto, methylenedioxy, ethylenedioxy or optionally substituted phenyl;


the term "heteroalkyl" refers to an C<sub>1-6</sub> alkyl radical wherein one, two or three hydrogen atoms have been replaced with a substituent independently selected from the group  
 30 consisting of -OR<sup>a</sup>, -NR<sup>b</sup>R<sup>c</sup>, and -S(O)<sub>n</sub>R<sup>d</sup> (where n is an integer from 0 to 2), with the understanding that the point of attachment of the heteroalkyl radical is through a carbon atom, wherein R<sup>a</sup> is hydrogen, acyl, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, or C<sub>3-7</sub> cycloalkyl C<sub>1-6</sub> alkyl; R<sup>b</sup> and R<sup>c</sup> are independently of each other hydrogen, acyl, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, or C<sub>3-7</sub>

- 86 -

cycloalkyl C<sub>1-6</sub> alkyl; when n is 0, R<sup>d</sup> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, or C<sub>3-7</sub> cycloalkyl C<sub>1-6</sub> alkyl, and when n is 1 or 2, R<sup>d</sup> is C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkyl C<sub>1-6</sub> alkyl, amino, acylamino, mono C<sub>1-6</sub> alkylamino, or di C<sub>1-6</sub> alkylamino;

the term "heterocyclyl" refers to a saturated or unsaturated non-aromatic cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from NR<sup>x</sup> {wherein each R<sup>x</sup> is independently hydrogen, C<sub>1-6</sub> alkyl, acyl, C<sub>1-6</sub> alkylsulfonyl, aminosulfonyl, (C<sub>1-6</sub> alkylamino)sulfonyl, (di C<sub>1-6</sub> alkylamino)sulfonyl, carbamoyl, (C<sub>1-6</sub> alkylamino)carbonyl, (di C<sub>1-6</sub> alkylamino)carbonyl, (carbamoyl) C<sub>1-6</sub> alkyl, (C<sub>1-6</sub> alkylamino)carbonyl C<sub>1-6</sub> alkyl, or di C<sub>1-6</sub> alkylaminocarbonyl C<sub>1-6</sub> alkyl}, O, or S(O)<sub>n</sub> (where n is an integer from 0 to 2), the remaining ring atoms being C and the heterocyclyl ring may be optionally substituted independently with one, two, or three substituents selected from C<sub>1-6</sub> alkyl, halo C<sub>1-6</sub> alkyl, heteroalkyl, halo, nitro, cyano C<sub>1-6</sub> alkyl, hydroxy, C<sub>1-6</sub> alkoxy, amino, mono C<sub>1-6</sub> alkylamino, di C<sub>1-6</sub> alkylamino, aryl C<sub>1-6</sub> alkyl, -(X)<sub>n</sub>-C(O)R (where X is O or NR', n is 0 or 1, R is hydrogen, C<sub>1-6</sub> alkyl, halo C<sub>1-6</sub> alkyl, hydroxy, C<sub>1-6</sub> alkoxy, amino, mono C<sub>1-6</sub> alkylamino, di C<sub>1-6</sub> alkylamino or optionally substituted phenyl, and R' is hydrogen or C<sub>1-6</sub> alkyl), -C<sub>1-6</sub> alkylene-C(O)R (where R is hydrogen, C<sub>1-6</sub> alkyl, halo C<sub>1-6</sub> alkyl, hydroxy, C<sub>1-6</sub> alkoxy, amino, mono C<sub>1-6</sub> alkylamino, di C<sub>1-6</sub> alkylamino or optionally substituted phenyl) or -S(O)<sub>n</sub>R<sup>d</sup> (where n is an integer from 0 to 2, and R<sup>d</sup> is hydrogen (provided that n is 0), C<sub>1-6</sub> alkyl, halo C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkyl C<sub>1-6</sub> alkyl, amino, mono C<sub>1-6</sub> alkylamino, di C<sub>1-6</sub> alkylamino, or hydroxy C<sub>1-6</sub> alkyl).

2. The compound according to claim 1, which is a compound of Formula (II):




(II)

wherein R<sup>1</sup>-R<sup>5</sup>, A, L, and X are as defined in claim 1.

3. The compound according to claim 1, which is a compound of Formula (III):


- 87 -



(III)

wherein R<sup>1</sup>-R<sup>5</sup>, A, L, and X are as defined in claim 1.

4. The compound according to any one of claims 1-3 wherein R<sup>1</sup> is methylene.
5. The compound according to any one of claims 1-3 wherein R<sup>2</sup> is 4-chlorophenyl or 3,4-dichlorophenyl.
6. The compound according to any one of claims 1-3 wherein R<sup>3</sup> is hydrogen.
7. The compound according to any one of claims 1-3 wherein L is -C(=O)-, -SO<sub>2</sub>-, -C(=O)N(R<sub>a</sub>)-, -C(=S)N(R<sub>a</sub>)-, or -C(=O)O-.
8. The compound according to claim 7 wherein L is -C(=O)-.
9. The compound according to claim 1, which is a compound of Formula (IV):




(IV)

wherein R<sup>3</sup>, R<sup>4</sup>, A, L, and X are as defined in claim 1.

10. The compound according to claim 1, which is a compound of Formula (VI):



- 88 -



(VI)


wherein X and R<sup>4</sup> are as defined in claim 1.

11. The compound according to claim 7 wherein X is absent, methylene, 1,2-ethanediyl,  
 5 1,3-propanediyl, or 1,4-butanediyl.

12. The compound according to claim 11 wherein R<sup>4</sup> is 3,4-dichlorophenyl,  
 3,4,5-trimethoxyphenyl, 4-methanesulfonylphenyl, 3-methanesulfonylphenyl,  
 4-methoxynaphthalen-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, phenyl,  
 3-fluorophenyl, 4-ethylphenyl, 3-methoxyphenyl, 2,4-difluorophenyl,  
 10 3-trifluoromethylphenyl, 4-methylphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-  
 carbamoylphenyl, 3-carbamoylphenyl, 4-acetylphenyl, 4-nitrophenyl, 2-methylphenyl, 2-  
 chloro-4-fluorophenyl, 3,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 2,3-dichlorophenyl,  
 2,4-dichlorophenyl, 4-bromophenyl, 4-chloro-3-nitrophenyl, 2-nitrophenyl, 2-nitro-4-  
 trifluoromethylphenyl, 4-chlorophenyl, 3-chlorophenyl, 2-chlorophenyl, 3-methylphenyl,  
 15 2-trifluoromethylphenyl, 2-methoxyphenyl, 3-bromophenyl, 4-trifluoromethylphenyl, 3-  
 trifluoromethylphenyl, 3,5-bis-trifluoromethylphenyl, 4-*tert*-butylphenyl, 4-ethoxyphenyl,  
 3-cyanophenyl, 4-cyanophenyl, 4-methoxyphenyl, 3-nitrophenyl, 3,5-dimethoxyphenyl, 4-  
 iodophenyl, 4-isopropylphenyl, 3-methoxycarbonylphenyl, 3-acetylphenyl, 2-  
 methylphenyl, indol-2-yl, 5-methoxyindol-2-yl, 5-chloroindol-2-yl, 2-  
 20 methoxycarbonylphenyl, 3,5-dichlorophenyl, 1-naphthyl, 3-chloro-2-methylphenyl, 2,5-  
 dimethylphenyl, 2-thienyl, 3-ethoxyphenyl, 3-isoquinolyl, 2-methylquinolin-6-yl, 3-  
 methylaminophenyl, 3-quinolyl, 2-quinolyl, 5-hydroxynaphthalen-2-yl, 8-  
 hydroxyquinolin-2-yl, 5,7-dimethyl-[1,8]naphthyridin-2-yl, 6-quinolyl, 3-  
 (acetylamino)phenyl, or 2,3,4-trimethoxyphenyl.


25 13. The compound according to claim 12 wherein R<sup>4</sup> is 3,4,5-trimethoxyphenyl, 4-  
 acetyl-phenyl, 3- carbamoylphenyl, 4-carbamoylphenyl, 3-methanesulfonylphenyl or 4-  
 methanesulfonyl-phenyl.

14. The compound according to claim 8, wherein X is  $-\text{CH}_2\text{S}-$  and  $\text{R}^4$  is 5-(3,4-dimethoxyphenyl)-pyrimidin-2-yl, 5-(3,4-methylenedioxyphenyl)-pyrimidin-2-yl or 5-(4-methoxyphenyl)pyrimidin-2-yl, and a salt thereof.
15. The compound according to claim 1, wherein ring A is  $\text{C}_{3-7}$  cycloalkyl or heterocycl  
5 or substituted phenyl.
16. The compound according to claim 15, wherein ring A is  $\text{C}_{3-7}$  cycloalkyl or heterocycl.
17. The compound according to claim 1, wherein  $\text{R}^2$  is substituted phenyl.
18. A method for preparing a compound of Formula (I) according to claim 1, wherein L  
10 is  $-\text{C}(=\text{O})\text{NR}_a-$ ,  $\text{R}_a$  is hydrogen, comprising reacting a compound of Formula (Ia)



with an isocyanate of a formula;  $\text{R}^4-\text{N}=\text{C}=\text{O}$ .

19. A method for preparing a compound of Formula (I) according to claim 1, wherein L is  $-\text{C}(=\text{O})-$ , comprising reacting a compound of Formula (Ia)



with a compound of a formula;  $\text{R}^4-\text{C}(=\text{O})\text{OH}$ .

20. A composition containing a therapeutically effective amount of a compound according to any one of claims 1-17, or a salt thereof; and an excipient.
21. A compound according to any one of claims 1-17, or a salt thereof for use in medical therapy or diagnosis.
- 5 22. A use of a compound of Formula (I) according to any one of claims 1-17, or a salt thereof; for the manufacture of a medicament comprising one or more compounds according to any one of claims 1-17, or a salt thereof for the treatment of a disease treatable by a CCR-3 receptor antagonist.
23. The use according to claim 22, wherein the disease is asthma.
- 10 24. The invention as herein before described, particularly with reference to the new compounds, intermediates, medicaments, uses and processes.

\*\*\*

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/13218

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C07D401/12 C07D211/18 C07D409/12 C07D401/04 C07D409/04 A61K31/454 A61P11/06		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, CHEM ABS Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 35452 A (DU PONT PHARM CO) 22 June 2000 (2000-06-22) abstract Core a,c table 3.1 claims 1-3 -----	1-24
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family		
Date of the actual completion of the international search 24 February 2003		Date of mailing of the international search report 05/03/2003
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Diederens, J

# INTERNATIONAL SEARCH REPORT

patent family members

International Application No

PCT/EP 02/13218

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0035452	A	22-06-2000	
		AU 1940600 A	03-07-2000
		AU 2057200 A	03-07-2000
		AU 2482100 A	03-07-2000
		AU 3126600 A	03-07-2000
		AU 3126700 A	03-07-2000
		BR 9917038 A	02-04-2002
		CN 1335771 T	13-02-2002
		EP 1161240 A1	12-12-2001
		EP 1140086 A1	10-10-2001
		EP 1158980 A1	05-12-2001
		EP 1140087 A1	10-10-2001
		EP 1156807 A1	28-11-2001
		JP 2002532427 T	02-10-2002
		NO 20012977 A	20-08-2001
		TR 200101859 T2	21-12-2001
		US 2003013741 A1	16-01-2003
		WO 0035449 A1	22-06-2000
		WO 0035451 A1	22-06-2000
		WO 0035452 A1	22-06-2000
		WO 0035453 A1	22-06-2000
		WO 0035454 A1	22-06-2000
		US 6492400 B1	10-12-2002
		US 6331541 B1	18-12-2001
		US 6486180 B1	26-11-2002
		US 6444686 B1	03-09-2002
		ZA 200103756 A	09-05-2002

**DERWENT-ACC-NO:** 2003-577220

**DERWENT-WEEK:** 200770

*COPYRIGHT 2008 DERWENT INFORMATION LTD*

**TITLE:** New piperidine derivatives useful in the treatment of chemokine receptor-3 mediated diseases e.g. asthma, psoriasis, and dermatitis

**INVENTOR:** BEIHAN W; BERNARD S D ; BRIAN S E ; DU BOIS D J ; JOE D B D ; JOHN K D ; KERTESZ D J ; SJOGREN E B ; SMITH D B ; WANG B

**PATENT-ASSIGNEE:** HOFFMANN LA ROCHE & CO AG F[HOFF] ,  
HOFFMANN LA ROCHE CO LTD F[HOFF]

**PRIORITY-DATA:** 2001US-334653P (November 30, 2001)

**PATENT-FAMILY:**

<b>PUB-NO</b>	<b>PUB-DATE</b>	<b>LANGUAGE</b>
WO 03045937 A1	June 5, 2003	EN
AU 2002352123 A1	June 10, 2003	EN
EP 1453825 A1	September 8, 2004	EN
BR 200214613 A	September 14, 2004	PT
KR 2004062665 A	July 7, 2004	KO
JP 2005515193 W	May 26, 2005	JA
CN 1599733 A	March 23, 2005	ZH
MX 2004005176 A1	September 1, 2004	ES
CN 1286831 C	November 29, 2006	ZH
MX 244991 B	April 13, 2007	ES

**DESIGNATED-STATES:** AE AG AL AM AT AU AZ BA BB BG BR BY  
 BZ CA CH CN CO CR CU CZ DE DK DM DZ  
 EC EE ES FI GB GD GE GH GM HR HU ID  
 IL IN IS JP KE KG KP KR KZ LC LK LR  
 LS LT LU LV MA MD MG MK MN MW MX MZ  
 NO NZ OM PH PL PT RO RU SD SE S G  
 SI SK SL TJ TM TN TR TT TZ UA UG UZ  
 VN YU ZA ZM ZW AT BE BG CH CY CZ DE  
 DK EA EE ES FI FR GB GH GM GR IE IT  
 KE LS LU MC MW MZ NL OA PT SD SE SK  
 SL SZ TR TZ UG ZM ZW AL AT BE BG CH  
 CY CZ DE DK EE ES FI FR GB GR IE IT  
 LI LT LU LV MC MK NL PT RO SE SI SK  
 T R

**APPLICATION-DATA:**

<b>PUB-NO</b>	<b>APPL- DESCRIPTOR</b>	<b>APPL-NO</b>	<b>APPL-DATE</b>
WO2003045937A1	N/A	2002WO- EP13218	November 25, 2002
AU2002352123A1	N/A	2002AU- 352123	November 25, 2002
BR 200214613A	N/A	2002BR- 014613	November 25, 2002
CN 1599733A	N/A	2002CN- 823978	November 25, 2002
CN 1286831C	N/A	2002CN- 823978	November 25, 2002
EP 1453825A1	N/A	2002EP- 787796	November 25, 2002
EP 1453825A1	N/A	2002WO- EP13218	November 25, 2002
BR 200214613A	N/A	2002WO- EP13218	November 25, 2002
JP2005515193W	N/A	2002WO- EP13218	November 25, 2002

MX2004005176A1	N/A	2002WO- EP13218	November 25, 2002
MX 244991B	N/A	2002WO- EP13218	November 25, 2002
JP2005515193W	N/A	2003JP- 547387	November 25, 2002
MX2004005176A1	N/A	2004MX- 005176	May 28, 2004
MX 244991B	N/A	2004MX- 005176	May 28, 2004
KR2004062665A	Based on	2004KR- 708357	May 31, 2004

**INT-CL-CURRENT:**

<b>TYPE</b>	<b>IPC DATE</b>
CIPP	A61K31/445 20060101
CIPP	C07D401/12 20060101
CIPS	A61K31/445 20060101
CIPS	A61K31/4525 20060101
CIPS	A61K31/4535 20060101
CIPS	A61K31/454 20060101
CIPS	A61K31/4545 20060101
CIPS	A61K31/4709 20060101
CIPS	A61K31/4725 20060101
CIPS	A61K31/506 20060101
CIPS	A61P11/06 20060101
CIPS	A61P43/00 20060101
CIPS	C07D211/16 20060101
CIPS	C07D211/18 20060101
CIPS	C07D401/04 20060101
CIPS	C07D401/12 20060101
CIPS	C07D405/04 20060101



CIPS C07D409/04 20060101  
CIPS C07D409/12 20060101  
CIPS C07D471/04 20060101

**RELATED-ACC-NO:** 2003-532727 2003-586792

**ABSTRACTED-PUB-NO:** WO 03045937 A1

**BASIC-ABSTRACT:**

NOVELTY - Piperidine derivatives (I), their prodrugs, isomers and salts are new.

DESCRIPTION - Piperidine derivatives of formula (I), or their prodrugs, individual isomers, racemic and non-racemic mixture of isomers, or salts are new.

R1 = 1-2C alkylene;

R2 = T1;

T1 = phenyl (optionally substituted by at least one T2);

T2 = 1-6C alkyl, halo-(1-6C)alkyl, OH-(1-6C)alkyl, Q1, C(O)R, C(O)R-amino, amino, (di)-1-6C alkylamino, 1-6C alkylthio, 1-6C alkylsulfinyl, 1-6C alkylsulfonyl, SO<sub>2</sub>NRcRd, 1-6C alkoxy, halo-(1-6C)alkoxy, 1-6C alkoxy carbonyl, carbamoyl, OH, halo, nitro, cyano, mercapto, methylenedioxy or ethylenedioxy;

R3 = H, 1-6C alkyl, C(O)R, Q', or Q'-(1-6C) alkyl;

A = 3-7C cycloalkyl, Q or T1;

L = -C(=O)-, -C(=S)-, -SO<sub>2</sub>, -C(=O)N(Ra)-, -C(=S)N(Ra)-, -SO<sub>2</sub>N(Ra)-, -C(=O)O-, -C(=S)O- or S(=O)<sub>2</sub>O-;

R4 = Q' or Q2;

Ra = H, 1-6C alkyl, C(O)R, Q', Q'(1-6C)alkyl, 1-6C alkoxy carbonyl or benzyloxy carbonyl;

R5, Rb, Rc, Rd, Rf, Rg and Rh = H or 1-6C alkyl;

X = absent, -C(RgRh)O, -C(RgRh)S-, -C(RgRh)NRb or 1-6C alkylene;

R = H, 1-6C alkyl, 3-7C cycloalkyl, 3-7C cycloalkyl-(1-6C)alkyl, phenyl or phenyl-(1-6C)alkyl;

Q = optionally saturated non-aromatic cyclic radical of 3 - 8 ring atoms in which at least two ring atoms are NR<sub>x</sub>, O or S(O)<sub>n</sub>, and the remaining ring atoms are C (optionally mono- - tri-substituted by (halo)1-6C alkyl, Q<sub>1</sub>, halo, nitro, cyano-(1-6C)alkyl, OH, 1-6C alkoxy, amino, mono-(1-6C)alkylamino, di(1-6C)alkylamino, Q'-(1-6C)alkyl, -(X)<sub>n'</sub>-C(O)Re, 1-6C alkylene-C(O)Re or -S(O)<sub>n</sub>R'd);

Q' = monocyclic or bicyclic aromatic hydrocarbon radical of 6 - 10 ring atoms (optionally substituted by at least one T<sub>2</sub>);

Q<sub>2</sub> = monocyclic or bicyclic radical of 5 - 12 ring atoms having at least one aromatic ring containing 1 - 3 N, O or S and the remaining ring atoms are C (optionally substituted by at least one T<sub>2</sub> or T<sub>1</sub>);

Q<sub>1</sub> = 1-6C alkyl radical (in which 1 - 3 H atoms are replaced by -OR'a, -NRb<sub>1</sub>Rc<sub>1</sub> or -S(O)<sub>n</sub>Rd<sub>1</sub>);

X = O or NR<sub>f</sub>;

Re = H, (halo)1-6C alkyl, OH, 1-6C alkoxy, amino, mono-(1-6C)alkylamino, di-(1-6C)alkylamino or T<sub>1</sub>;

n = 0 - 2;

$n' = 0$  or  $1$ ;

$R'd = H$ , (halo)-1-6C alkyl, 3-7C cycloalkyl, 3-7C cycloalkyl-(1-6C) alkyl, amino, mono-(1-6C)alkylamino, di-(1-6C) alkylamino or OH-(1-6C) alkyl;

$R_x = H$ , 1-6C alkyl,  $C(O)R$ , 1-6C alkylsulfonyl, aminosulfonyl, (di)-1-6C alkylaminosulfonyl, carbamoyl, (di)-1-6C alkylaminocarbonyl, (carbamoyl)-(1-6C)alkyl or (di)-1-6C (alkylamino)carbonyl-(1-6C)alkyl;

$R'a$ ,  $R_{b1}$  and  $R_{c1} = H$ ,  $C(O)R$ , 1-6C alkyl, 3-7C cycloalkyl or 3-7C cycloalkyl-(1-6C)alkyl;

$R_{d1} = H$ , 1-6C alkyl, 3-7C cycloalkyl, 3-7C cycloalkyl-(1-6C)alkyl, amino,  $C(O)R$ -amino, mono-(1-6C)alkylamino or di-(1-6C)alkylamino.

$Q1$  is attached through a carbon atom.

Provided that:

(1) when  $R_1$  is  $-CH_2-$ ,  $R_2$  and  $A$  are phenyl,  $R_3$  and  $R_5$  are  $H$ ,  $L$  is  $-C(=O)NH-$  and  $X$  is absent, then  $R_4$  is other than 2,4-difluorophenyl;

(2) when  $n$  is  $0$ , then  $R_{d1}$  is  $H$ , 1-6C alkyl, 3-7C cycloalkyl, or 3-7C cycloalkyl-(1-6C)alkyl;

(3) when  $n$  is  $1$  or  $2$ , then  $R_{d1}$  is 1-6C alkyl, 3-7C cycloalkyl, 3-7C cycloalkyl-(1-6C)alkyl, amino,  $C(O)R$ -amino, mono-(1-6C)alkylamino, or di-(1-6C)alkylamino; and

(4) when  $n$  is  $0$ , then  $R'd$  is  $H$ .

An INDEPENDENT CLAIM is also included for the preparation of (I).

## Chemokine (CCR-3) Antagonist.

The CCR-3 antagonistic activity of N-(3-(3-(2-(4-(4-chlorobenzyl)-piperidin-1-yl)-cyclopentyl)-ureido)-phenyl)-acetamide (Ia) was determined by its ability to inhibit the binding of I125-eotaxin to CCR-3 L1.2 transfectant cells (see Ponath, P.D. et al., J. Clin. Invest. 97:604-612 (1996)). (Ia) showed an IC50 value of 0.0185 microM.

USE - (I) is used in the manufacture of a medicament for the treatment of a disease treatable by a chemokine (CCR-3) receptor antagonist (e.g. asthma) (claimed). Also useful for treating eosinophil-induced diseases including inflammatory or allergic diseases, respiratory allergic diseases (e.g. allergic rhinitis, hypersensitivity lung disease, hypersensitivity pneumonitis, eosinophilic pneumonias, (e.g. chronic eosinophilic pneumonia), inflammatory bowel disease (e.g. Crohn's disease, and ulcerative colitis), psoriasis, and dermatoses (e.g. dermatitis and eczema).

ADVANTAGE - (I) are potent chemokine (CCR-3) receptor antagonists. (I) inhibit the binding of eotaxin to the CCR-3 receptor, hence facilitate treatment of eosinophil induced disease.

**EQUIVALENT-ABSTRACTS:**

## ORGANIC CHEMISTRY

Preparation (claimed): Preparation of (I) comprises either:

(a) process A: Preparation of (I) (where L is -C(=O)NRa; and Ra is H) involves reacting a piperidine derivative of formula (Ia) with an isocyanate of formula R4-N=C=O; or

(b) Process B: Preparation of (I) (where L is -C(=O))

involves reacting (Ia) with an acid of formula  $R_4-C(=O)OH$ .

#### Preferred Definitions:

$R_1$  = methylene;

$R_2$  = 4-chlorophenyl, or 3,4-dichlorophenyl;

$R_3$  = H;

$L$  =  $-C(=O)-$ ;

$X$  = absent,  $CH_2S$ , methylene, 1,2-ethanediyl, 1,3-propanediyl, or 1,4-butanediyl;

$R_4$  = 5-(3,4-dimethoxyphenyl)-pyrimidin-2-yl, 5-(3,4-methylenedioxyphenyl)-pyrimidin-2-yl, 5-(4-methoxyphenyl)-pyrimidin-2-yl, 3,4,5-trimethoxyphenyl, 4-acetyl-phenyl, 3-carbamoylphenyl, 4-carbamoylphenyl, 3-methanesulfonylphenyl, or 4-methanesulfonyl-phenyl; and

$A$  = 3-7C cycloalkyl or heterocyclyl.

An intermediate of formula (Ia) has been disclosed to be new.

The dosage is 0.01 - 20 (preferably 0.1 - 10) mg/kg. The route of administration is oral, transdermal, inhalation (e.g. intranasal or oral inhalation), or parenteral (e.g. intramuscular, intravenous, or subcutaneous).

#### SPECIFIC COMPOUNDS

10 Compounds (I) are specifically disclosed, e.g. N-(3-(3-(2-(4-(4-Chlorobenzyl)-piperidin-1-yl)-cyclopentyl)-ureido)-phenyl)-acetamide (Ia).

4-(4-chlorobenzyl)piperidine (52 mg) was alkylated with 7-oxabicyclo(4.1.0)heptane (0.25 ml) in ethanol (0.5 ml) at 80 degrees C for 3 days to give (+/-)-trans-2-(4-(4-chlorobenzyl)piperidin-1-yl)cyclohexanol (A) (68 mg).

(A) (390 mg) in dichloromethane (6 ml) at 0 degrees C was treated successively with triethylamine (350 microl) and MeSO<sub>2</sub>Cl (194 microl), stirred for 2 hours at 0 degrees C and partitioned between dichloromethane and ammonium hydroxide. A solution of the residue in tetrahydrofuran (3 ml) and 28-30% ammonium hydroxide (1.2 ml) was stirred for 24 hours at 70 degrees C, allowed to cool to room temperature and partitioned between ethyl acetate and 1N sodium hydroxide. Work up gave (+/-)-trans-2-(4-(4-Chlorobenzyl)piperidin-1-yl)-cyclohexylamine (B) (260 mg).

(B) (56 mg) and 5-isocyanato-1,2,3-trimethoxybenzene (76 mg) were coupled in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) in dimethylformamide (1 ml) at room temperature for 1.5 hour. After work up (+/-)-trans-1-(2-(4-(4-chlorobenzyl)piperidin-1-yl)-cyclohexyl)-3-(3,4,5-trimethoxyphenyl)urea (Ia) (52 mg) was obtained.

**TITLE-TERMS:** NEW PIPERIDINE DERIVATIVE USEFUL TREAT  
RECEPTOR MEDIATOR DISEASE ASTHMA  
PSORIASIS DERMATITIS

**DERWENT-CLASS:** B02 B03

**CPI-CODES:** B06-H; B07-D05; B14-C03; B14-E08; B14-E10C; B14-G02A; B14-K01; B14-L06; B14-N04; B14-N17C;

**CHEMICAL--CODES:** Chemical Indexing M2 \*01\*  
Fragmentation Code F011 F014 F433 G013  
G019 G033 G112 G553 H1 H161 H2 H201 H5  
H541 H6 H602 H641 H8 K0 L4 L420 M1  
M123 M129 M132 M137 M210 M211 M272  
M281 M311 M321 M342 M413 M510 M521  
M532 M541 M710 P420 P431 P714 P738  
P820 P822 P831 P924 P943 Specific  
Compounds RAB53B Registry Numbers  
751707

Chemical Indexing M2 \*02\*  
Fragmentation Code C116 C216 C316 D010  
D019 D020 D021 D022 D029 D040 D049  
D140 D150 F010 F011 F014 F017 F019  
F020 F021 F433 G001 G002 G003 G010  
G011 G012 G013 G014 G015 G016 G019  
G020 G021 G022 G029 G030 G039 G040  
G050 G100 G111 G112 G113 G221 G299  
G553 G563 H1 H100 H101 H102 H103 H121  
H122 H123 H141 H142 H143 H161 H181  
H182 H183 H2 H201 H321 H322 H323 H341  
H342 H343 H401 H402 H403 H404 H405  
H421 H422 H423 H424 H441 H442 H443  
H444 H481 H482 H483 H484 H492 H494  
H498 H521 H522 H523 H541 H542 H543  
H592 H594 H599 H600 H608 H609 H621  
H622 H623 H641 H642 H643 H681 H682  
H683 H684 H685 H686 H689 J011 J012  
J013 J014 J111 J112 J113 J171 J172  
J173 J211 J212 J221 J222 J231 J232  
J271 J272 J273 J311 J312 J321 J322  
J331 J332 J341 J342 J351 J361 J371  
J372 J373 J390 J411 J412 J431 J432  
J471 J472 J521 J581 J582 J583 K340  
K351 K352 K353 K399 K442 K499 L142  
L143 L145 L199 L410 L420 L431 L432  
L450 L462 L463 L472 L499 L531 L532  
L599 L640 L650 L660 L699 M1 M113 M116  
M121 M122 M123 M124 M125 M126 M129

M131 M132 M135 M136 M137 M139 M142  
M143 M147 M148 M149 M150 M210 M211  
M212 M213 M214 M215 M216 M220 M221  
M222 M223 M224 M225 M226 M231 M232  
M233 M240 M262 M271 M272 M273 M280  
M281 M282 M283 M311 M312 M313 M314  
M315 M316 M321 M322 M323 M331 M332  
M333 M334 M340 M342 M343 M344 M349  
M353 M362 M372 M373 M381 M383 M391  
M392 M393 M412 M413 M510 M511 M512  
M513 M521 M522 M523 M530 M531 M532  
M533 M540 M541 M542 M543 M710 P420  
P431 P714 P738 P820 P822 P831 P924  
P943 Ring Index Numbers 01662 Markush  
Compounds 009880401

Chemical Indexing M2 \*03\*

Fragmentation Code F011 F014 F433 G012  
G013 G033 G112 G553 H1 H161 H2 H201 H6  
H602 H641 J0 J011 J3 J341 K0 L4 L432  
M1 M123 M129 M132 M137 M210 M211 M262  
M281 M311 M321 M342 M413 M510 M521  
M532 M541 M710 M720 N234 N261 N331  
N361 N512 P420 P431 P714 P738 P820  
P822 P831 P924 P943 Specific Compounds  
RAB52G Registry Numbers 751675

Chemical Indexing M2 \*04\*

Fragmentation Code D012 D621 F011 F014  
F433 G013 G033 G111 G553 H1 H161 H2  
H201 H6 H602 H641 K0 L4 L432 L943 M1  
M123 M126 M132 M137 M280 M311 M321  
M342 M412 M511 M521 M531 M541 M710  
M720 N234 N261 N331 N361 N512 P420  
P431 P714 P738 P820 P822 P831 P924  
P943 Specific Compounds RAB52M  
Registry Numbers 751682

Chemical Indexing M2 \*05\*

Fragmentation Code F011 F012 F014 F211



F433 G013 G033 G111 G553 H1 H161 H2  
H201 H6 H602 H641 K0 L4 L432 M1 M123  
M132 M280 M311 M312 M321 M332 M342  
M373 M391 M413 M510 M522 M531 M541  
M710 M720 N234 N261 N331 N361 N512  
P420 P431 P714 P738 P820 P822 P831  
P924 P943 Specific Compounds RAB52Q  
Registry Numbers 751686

## Chemical Indexing M2 \*06\*

Fragmentation Code F011 F014 F433 G013  
G017 G033 G112 G563 H1 H161 H2 H201 H5  
H543 H6 H602 H641 H8 K0 L4 L432 M1  
M123 M129 M132 M137 M210 M211 M272  
M283 M311 M321 M342 M413 M510 M521  
M532 M541 M710 M720 N234 N261 N331  
N361 N512 P420 P431 P714 P738 P820  
P822 P831 P924 P943 Specific Compounds  
RAB52Z Registry Numbers 751695

## Chemical Indexing M2 \*07\*

Fragmentation Code C116 C216 C316 D010  
D019 D020 D021 D022 D029 D040 D049  
D140 D150 F010 F011 F014 F017 F019  
F020 F021 F433 G001 G002 G003 G010  
G011 G012 G013 G014 G015 G016 G019  
G020 G021 G022 G029 G030 G039 G040  
G050 G100 G111 G112 G113 G221 G299  
G553 G563 H1 H100 H101 H102 H103 H121  
H122 H123 H141 H142 H143 H161 H181  
H182 H183 H2 H201 H321 H322 H323 H341  
H342 H343 H401 H402 H403 H404 H405  
H421 H422 H423 H424 H441 H442 H443  
H444 H481 H482 H483 H484 H492 H494  
H498 H521 H522 H523 H541 H542 H543  
H592 H594 H599 H600 H608 H609 H621  
H622 H623 H641 H642 H643 H681 H682  
H683 H684 H685 H686 H689 J011 J012  
J013 J014 J111 J112 J113 J171 J172  
J173 J211 J212 J221 J222 J231 J232

J271 J272 J273 J311 J312 J321 J322  
J331 J332 J341 J342 J371 J372 J373  
J411 J412 J431 J432 J471 J472 J521  
J581 J582 J583 K0 K353 K399 K442 K499  
L142 L143 L145 L199 L4 L431 L432 L463  
L472 L499 L532 L640 L650 L699 M1 M113  
M116 M121 M122 M123 M124 M125 M126  
M129 M131 M132 M135 M136 M137 M139  
M142 M143 M149 M150 M210 M211 M212  
M213 M214 M215 M216 M220 M221 M222  
M223 M224 M225 M226 M231 M232 M233  
M240 M262 M271 M272 M273 M280 M281  
M282 M283 M311 M312 M313 M314 M315  
M316 M321 M322 M323 M331 M332 M333  
M334 M340 M342 M343 M344 M349 M353  
M362 M372 M373 M383 M391 M392 M393  
M412 M413 M510 M511 M512 M513 M521  
M522 M523 M530 M531 M532 M533 M540  
M541 M542 M543 M710 M720 N234 N261  
N331 N361 N512 P420 P431 P714 P738  
P820 P822 P831 P924 P943 Ring Index  
Numbers 01662 Markush Compounds  
009880402

## Chemical Indexing M2 \*08\*

Fragmentation Code C116 C216 C316 D010  
D019 D020 D021 D022 D029 D040 D049  
D140 D150 F010 F011 F014 F017 F019  
F020 F021 F433 G001 G002 G003 G010  
G011 G012 G013 G014 G015 G016 G019  
G020 G021 G022 G029 G030 G039 G040  
G050 G100 G111 G112 G113 G221 G299  
G553 G563 H1 H100 H101 H102 H103 H121  
H122 H123 H141 H142 H143 H161 H181  
H182 H183 H2 H201 H321 H322 H323 H341  
H342 H343 H401 H402 H403 H404 H405  
H421 H422 H423 H424 H441 H442 H443  
H444 H481 H482 H483 H484 H492 H494  
H498 H521 H522 H523 H541 H542 H543  
H592 H594 H599 H600 H608 H609 H621

H622 H623 H641 H642 H643 H681 H682  
H683 H684 H685 H686 H689 J011 J012  
J013 J014 J111 J112 J113 J171 J172  
J173 J211 J212 J221 J222 J231 J232  
J271 J272 J273 J311 J312 J321 J322  
J331 J332 J341 J342 J361 J371 J372  
J373 J411 J412 J431 J432 J471 J472  
J521 J581 J582 J583 K353 K399 K442  
K499 L142 L143 L145 L199 L432 L463  
L472 L499 L532 L650 L699 M1 M113 M116  
M121 M122 M123 M124 M125 M126 M129  
M131 M132 M135 M136 M139 M142 M143  
M149 M150 M210 M211 M212 M213 M214  
M215 M216 M220 M221 M222 M223 M224  
M225 M226 M231 M232 M233 M240 M262  
M271 M272 M273 M280 M281 M282 M283  
M311 M312 M313 M314 M315 M316 M321  
M322 M323 M331 M332 M333 M334 M340  
M342 M343 M344 M349 M353 M362 M372  
M373 M381 M383 M391 M392 M393 M412  
M413 M510 M511 M512 M513 M521 M522  
M523 M530 M531 M532 M533 M540 M541  
M542 M543 M710 M720 N241 N261 N331  
P420 P431 P714 P738 P820 P822 P831  
P924 P943 Ring Index Numbers 01662  
Markush Compounds 009880403

**SECONDARY-ACC-NO:****CPI Secondary Accession Numbers:** 2003-156020